235 0 15

THE

BOTANIST'S CALENDAR,

AND

POCKET FLORA.



Glad ZEPHYR floats on light aurelian wing, Scatt'ring sweet florets in the lap of Spring; O'er Summer's bosom breathing fragrance pours, Rifl'ing gay FLORA of her honied stores.

Ephemendes



BOTANIST'S CALENDAR,

AND

POCKET FLORA:

ARRANGED ACCORDING TO THE

LINNÆAN SYSTEM.

To which are added,

REFERENCES TO THE BEST FIGURES

OF

BRITISH PLANTS.

IN TWO VOLUMES.

VOLUME I.

LONDON:

PRINTED BY T. BENSLEY,

FOR B. AND J. WHITE, FLEET-STREET.

1797.



Entered at Stationers' Hall.

PREFACE.

This Little Work is submitted to the public eye with the utmost dissidence. The Compiler is aware that the complete Botanist may object to the partial notice of the Grasses and Rushes, and to the total omission of the minute and numerous Tribes belonging to the Class Cryptogamia: but to have included these, would have carried him beyond the limits of a Pocket Compendium; which he now publishes, with the humble view only of leading the Student in Botany through the delightful wilds of blooming vegetation, to a systematic knowledge of those indigenous Plants

which are peculiarly striking, or beautiful: and to assist the investigating FAIR ONE, who, with nice and curious hand, crops a FLOWER-ING RIVAL of its blushing Sweets!

South Lambeth, Мау 30, 1797.

ABBREVIATIONS.

Those in the Margin, are the Classes and Orders.

With. Arr. - Withering's Arrangement of British Plants, (third Edition, 4 Vols.)

Curt. - - - Curtis's Flora Londinenfis, 6 vols.

Woodv. - - Woodville's Medical Botany, 4 vols.

E. Bot. - - English Botany, 6 vols.


Fl. Ruft. - - Flora Ruftica, 4 vols.

White's Nat. Cal. White's Naturalist's Calendar, I vol.

Gen. Ch. - Generic Character.

The Months abbreviated shew the continuance of flowering.

The figures of reference are to the Page in Withering's Arrangement; to the Plate in Woodville's Medical Botany, English Botany, and Flora Rustica; and to the Volume in Curtis's Flora Londinensis.



Botanist's Calendar.

JANUARY.

PRIMULA. Bloff. the tube cylindrical; mouth open: Stem within the tube: Capf. 1-celled, cylindrical, many-feeded, opening with 10 teeth: Summit a knob.

P. Leaves wrinkled, toothed; bor-vulgáris. der of the bloffom flat.

With. Arr. 233. Curt. 6. E. Bot. 4.

Common Primrofe. Woods, hedges,

thickets, and heaths, (particularly in a clayey foil.) May. Pen. Mo.

I have given the first place in this Epitomized Flora to

"The pale primrofe, that dies unmarried," on the authority of that curious observer of nature, the late

Reverend Gilbert White, of Selborne, whose taste and leifure led him to watch "The Loves of the Plants:" and to mark, with peculiar accuracy, the earliest and the latest periods of their flowering. man, who found amusement in every rural ftep he took:

"Tongues in the trees, books in the run-" ning brooks;

" Sermons in stones, and good in every " thing !"

In his Naturalists' Calendar he informs us, the Primrofe has flowered in mild winters on Nov. 10, and Jan. 6. In backward fprings, late as April 7.

GALA'NTHUS. Petals 3, concave: Nectary 3 smaller petals notched at the end: Summit undivided.

G. Bloff. white. Orchards. Meanivális. dows.

> With. Arr. 331. E. Bot. 19. Common Snow-drop. Fair Maids of February. March.

White's Nat. Cal. Dec.29—Feb.5.

Professor Martyn in Miller's im- Hex. Mo. proved Dictionary, January 25, 1796.

DA'PHNE, Calyx generally none. Bloff. 1 petal, regular, 4-cleft funnel-shaped: Drupa like a berry, 1-celled, superior.

D. Flowers fitting on the ftem Mezéreum. mostly 3 together: leaves spear-shaped, deciduous: berries glo-bular.

With. Arr. 376. Woodv. 23.

The terminating buds produce leaves; the central buds flowers. Red.

Mezereon. Spurge olive. Woods.

March. Oct. Mo.

White's Nat. Cal. Dec. 15-Feb. 15.

Helle'Borus. Bloff. none: Cal. 5 leaves, often coloured: Nectaries 2-lipped, tubular: Capf. like a legumen: many-feeded, rather upright, beaked.

).

ls

h.

fœ'tidus.

H. Stem many-flowered, leafy: leaves bird-footed.

Flowers numerous, somewhat globular, green, sometimes tinged with purple at the edges.

With Arr. 510. Woodv. 19.

Bearsfoot. Stinking Hellebore. Meadows, shady places, hedges.

Poly. Poly.

April.

White's Nat. Cal. Dec. 14-30.

vir idis.

H. Stem many-flowered, leafy; leaves finger-like.

Flowers mostly 2: yellowish green.

With. Arr. 510. Curt. 6. E. Bot. 200.

Poly. Poly.

Green flowered Hellebore. Woods, and dry pastures in chalky foil.

U'LEX. Calyx 2-leaved: Legumen fcarcely longer than the calyx.

europæ'us. U. Leaves woolly, acute; thorns fcattered.

Bloff. yellow, at the base of the thorns.

With, Arr. 626.

Gorze. Furze, Whins. Heaths, road fides, pastures. Aug. Diad. Dec. White's Nat. Cal. Dec. 4-Feb. 1.

VERO'NICA. Bloff. border 4-cleft; the lower fegment narrowest: Capf. 2 celled; notched at the end.

V. Flowers folitary: leaves heart- agres tis. shaped, on leaf-stalks, shorter than the fruit-stalks.

Curt. 1. With. Arr. 17.

Bloff. blue; fometimes white.

Germander. Procumbent Speedwell. Paftures, ploughed-fields.

April. Di. Mo.

White's Nat. Cal. Jan. 27-Mar. 15.

CRO'CUS. Bloff. with 6 equal divisions: Summits coiled.

C. Sheath 1 valve, rifing from the vérnus. root: tube of the bloffom very long. Summit in 3 fhort wedgeshaped lobes.

With. Arr. 68. E. Bot. 344.

Spring Suffron, or Crocus. Leaves broader, with flat edges.

White's Nat. Cal. Jan. 13-Mar. 18.

Tri, Mo.

Viola. Calyx 5-leaved, adhering to the blofform above the base: Bloff. 5 petals, irregular, with a spur behind: Anthers cohering: Caps. 1-celled, 3-valved.

fhaped, toothed: leaves eggfhaped, toothed: calyx fmooth,
but half the fize of the bloffom.
This plant is liable to almost endless varieties. Corn - fields,
ditch-banks.

With. Arr. 262. Curt. 1. Woodv. 252.

Pen. Mo. Pansies. Heart's-ease. Sept.. White's Nat. Cal. Jan. 3.

Lonice RA. Bloff. 1 petal tubular, irregular: Berry beneath; 1 to 3-celled; many-feeded.

Pericly'menum.

L. Heads egg-shaped, tiled, terminating: leaves distinct, deciduous: blossom gaping. Hedges.

With. Arr. 246. Curt. 1.

Leaves and stem smooth. Bloff. red on the outside, yellowish within.

Pen. Mo. Woodbine. Honeysuckle. June. White's Nat. Cal. Jan. 25.

Co'RYLUS. Male and female flowers on the fame plant: Bloff. none.

Male; Cal. 1 leaf, 3-cleft, refembling a fcale, containing 1 flower.

Fem. Cal. 2-leaved, ragged: Nut egg-shaped.

C. Stipulæ fpear-fhaped: twigs Avellána. hairy.

With. Arr. 378.

The styles being of a vivid crimfon have a very beautiful appearance.

Common hazel nut-tree. April.
Woods and hedges.

White's Nat. Cal. Jan. 3-Feb. 28. Oft. Di.

RANUN'CULUS. Cal. deciduous, 5-(or 3-) leaved: Petals 5 (rarely 2, 3, or 8), with a nectariferous scale or pore within the claw: Styles permanent: Seeds incrusted, upright. Aug.

R. Calyx expanding: fruit-stalks répens. furrowed: fuckers creeping: leaves compound.

Curt. 4. With. Arr. 509.

Poly. Poly. Creeping Crow-foot. Meadows. White's Nat. Cal. Jan. 13.

LA'MIUM. Bloff. upper lip entire, vaulted; lower-lip inversely heart-shaped: mouth with a bristle-shaped tooth on each side.

purpúreum. L. Leaves heart-shaped, blunt, on leaf-stalks.

With. Arr. 527. Curt. 1. Flowers 6 in each leaf, in a double row.

Red Dead-nettle. Archangel. Rubbish, corn-fields, gardens.

Didy. Gym.

Sept.

White's Nat. Cal. Jan. 3-21.

al'bum. L. Leaves heart-shaped, tapering to a point, ferrated, on leaf-stalks: flowers about 20 in a whirl.

With. Arr. 526. Curt. 2.

Flowers white; fometimes, though rarely, with a pinky tinge:

Anthers hairy, black.

White Archangel. White Deadnettle. Rubbish, corn-fields, ditch-banks. June.

Didy. Gym.

White's Nat. Cal. Jan. 13.

LEON'TODON. Receptacle naked: Calyx tiled; the inner scales parallel, equal: Down hair-like.

L. Outer scales of the calyx entire, reflected: leaves smooth,
notched, and acutely toothed.
With. Arr. 079. Curt. 1. Woodv. 7.
Bloff. yellow, expanding about 5
or 6 in the morning, and closing early in the afternoon.
Common Dandelion. Pissa-bed.

Meadows, road-fides. Sept. Syng. Æqua. White's Nat. Cal. Jan. 16—Mar. 11.

Sene cio. Recept. naked: Down hair-like, long: Calyx conical, double: the scales as if dead at the ends.

S. Leaves winged-indented, em- vulgáris. bracing the stem: flowers scattered.

With. Arr. 721. Curt. 1.

Bloffom yellow, terminating.

Common Groundfel. Simfon. Cultivated ground, rubbish. Sept. Syng Surer.

White's Nat. Cal. Jan. 3—15.

FEBRUARY.

ERIO'PHORUM. Hufks like chaff, tiled on every fide: Bloff. none: Seeds encompassed by very long wool-like hairs.

vaginátum. E. Straw cylindrical, sheathed: fpike skinny.

With. Arr. 71. Curt. 4.
Tri. Mo. Hare's-tail Rush. Single-headed
Cotton-grass. April.

Rus'cus. Stamens and piftils mostly on different plants: Bloff. none: Cal. 6 leaves: Nectury central, egg-shaped, open at the top.

Male flowers, Anthers united. Fem. flowers, Berry 3-celled: Seeds generally 2.

R. Leaves bearing the flowers on the upper furface, naked.

With. Arr. 68. Woodw. 237. Knee Holly. Butchers' Broom. Woods, thickets, hedges.

April.

aculeátus.

Tri. Mo.

BETULA. Male and female flowers on the same plant: Cal. 1 leaf, with 3 or 5 clefts: Bloff. with 4 divisions.

Male. Cal. 3-flowered.

Fem. Cal. 2-flowered: Seeds 2 or 3.

B. Fruit-stalks branched, wedge-al'nus. shaped, very blunt: leaves round-ish, glutinous: veins underneath woolly at the base.

With. Arr. 206.

Catkins brown. March. Tet. Di. Alder, Owler. Near water.

Vi'ola. See Gen. Ch. page 6.

V. Leaves heart-shaped: fuckers odorátz. creeping. Floral leaves above the middle of the fruit-stalk.

With. Arr. 260. Curt. 1.

Bloff. rich blue purple, fmelling very fweet.

Sweet Violet. April.

Warm hedge, and ditch-banks, Pen. Mo. moist warm lanes; in clay or marle.

FEBRUARY.

NARCIS'SUS. Bloff superior, of 6 equal petals: Nectary of 1 leaf, furrounding the stamens, mostly funnel-shaped.

pfuedonarcis sus. N. Sheath 1 flowered: nectary bell-shaped, upright, curled; as long as the petals: petals eggshaped.

With. Arr. 332. E. Bot. 17.

Stalk 2-edged. Petals egg-spear-shaped, straw-coloured. Nec. a full yellow: the rim a little plaited and snipt.

Hex. Mo.

Common Daffodil. March.
Woods, meadows, and fides of hedges.

CERA'STIUM. Calyx 5-leaved: Petals cloven: Capf. 1-celled, opening at the top.

púmilum.

C. Flowers with 5 stamens: petals cloven, as long as the calyx: capfule twice as long.

With. Arr. 435. Curt. 6.

Dec. Pen.

Mouse-Ear, March.
Dry banks near Croydon.

FRAGA'RIA. Calyx 10-cleft: Petals 5: Seeds naked, smooth, on a receptacle which is egg-shaped, coloured, deciduous, retembling a berry.

F. Stem proftrate, without creep- stérilis. ing runners.

With. Arr. 473. Curt. 3.

Flowers folitary white.

Barren Strawberry. May.

Barren pastures, heaths, hedgebanks.

White's Nat. Cal. Feb. 1-Mar. 26. Ico. Poly.

RANUN'CULUS. See Gen. Ch. page 7.

R. Leaves heart-shaped, angular, Ficária, on leaf-stalks: stem 1-flowered: flowers with 8 petals: calyx with 3 leaves.

With. Arr. 503. Curt. 2.

Petals 8 or 9, bright yellow.

Common Pilewort. Leffer Celandine. Meadows, pastures, Po'y. Poly. ditch-banks, South Lambeth.

LA'MIUM. See Gen. Ch. page 8.

amplexicaúle. L. Floral-leaves fitting, embracing the stem, blunt.

With. Arr. 527. Curt. 2.

Bloff. purple.

June.

Didy. Gym.

Great Henbit. Henbit Archangel.
Sandy corn-fields, cultivated grounds.

Tax'us. Male and female flowers on different plants: Bloff. none: Calyx a 4 or 7-leaved bud.

Male Anthers target-shaped, 8cleft.

Fem. Style none: Seed 1 furrounded at the base by a pulpy receptacle; the upper half naked.

baccáta.

T. Leaves folitary, ftrap-shaped, prickle-pointed, near together: receptacle of the male flowers fomewhat globular.

With. Arr. 614.

Berries when ripe, red. April.

Mon. Poly. Common Yew Tree.

White's Nat. Cal. Feb. 14 -- Mar. 27.

Tussila'Go. Recept. naked: Down hair-like: Calyx scales equal, as tall as the surface of the florets, somewhat membranaceous.

T. Stalk with 1 flower, tiled: Far'fara. leaves fomewhat heart-fhaped, angular, finely toothed.

With. Arr. 719. Curt. 2.

Bloff. yellow. April.

Common Coll's foot. Pastures and moistish places, in stiff-clayey Syng. Super. soil, and on limestone rubbish.

MARCH.

VERO'NICA. See Gen. Ch. page 5.

hederæfólia. V. Flowers folitary: leaves heartfhaped, flat, 5-lobed.

With. Arr. 17. Curt. 2.

Bloff. pale blue.

Small Henbit. Ivy-leaved Speedwell. Hedge-banks, ploughed fields.

FRAX'INUS. Cal. none, or with 4 divisions: Bloff. none, or with 4 petals.

Fem. and hermaphrodite flowers. Pift. 1. Capf. 2-celled; leaflike upwards, compressed; 1 cell barren: Seed spear-shaped.

F. Leafits ferrated: flowers without petals.

With. Arr. 57. Var. leaves fimple. T. H. W.

Gent. Mag. vol. iv. p. 598.

There is no place of growth of this curiofity mentioned, but

Di. Mo.

excel for.

the fenfible and intelligent reporter could not well be deceived about it. May. Di. Mo.

Common Ash Tree. Woods, hedgerows.

HIPPO'PHAE. Male and Fem. flowers on different plants: Bloff. none.

Male, Calyx 1 leaf, 2-lobed.

Fem. Calyx 1 leaf, tubular: Berry fuperior, 1-celled: Seed hard, fhining.

H. Leaves spear-shaped. With. Arr. 204. May.

Rhamnoides.

Sea Buckthorn. Common Sallow-thorn. Sea shore, in fand. Tet. Mo.

VI'OLA. See Gen. Ch. page 6.

V. Stem ascending as it attains its canina. full growth: leaves oblong heartshaped.

With. Arr. 261. Curt. 2.

Bloff. pale blue, streaked, scentlefs, fometimes all white, fometimes the fpur only white.

Dog's Violet. Tune. Pen. Mo.

Shady places, heaths, and hedgebanks.

White's Nat. Cal. Mar. 6—Ap.18. Var. 2. Dwarf. In all respects smaller.

Vi'ola. See Gen. Ch. page 6.

hir ta.

V. Leaves heart-shaped, rough, with strong hair. Leaf-stalks rough with hair: floral leaves below the middle of the fruit-stalk.

With. Arr. 260. Curt. 1.

Bloff. pale blue; scentless.

Hairy Violet. April.

Pen. Mo.

Shady places and hedges in a calcareous foil.

VIN'CA. Bloff. falver-shaped: Seedveff. upright little bags: Seeds naked.

minor.

V. Stem trailing: leaves fpearegg-shaped: flowers on fruitstalks.

With. Arr. 268. Curt. 3.

Pen. Mo.

Bloff. blue or white. May. Leffer Periwinkle. Woods and hedges. Rare. PULMONA'RIA. Bloff. funnel-shaped; mouth open: Calyx tubular, but 5-fided.

 P. Root-leaves egg-heart-shaped, officinális. rough; upper leaves egg-shaped, acute.

With. Arr. 228. E. Bot. 118. Woodv. 212.

Bloff. purplish red when newly expanded, foon changing to blue.

Spotted Lungwort. Cowslips of Jerusalem. May.

Cliff-wood, near Darlington. Pen Mo. White's Nat. Cal. Mar. 4—Ap. 16.

ULMUS. Cal. 5-cleft: Bloff. none: Capf. fuperior, 1-celled, leaf-like and compressed: Seed solitary.

U. Leaves doubly ferrated, unequal compétris. at the base: flowers almost sitting, crowded together.

With. Arr. 278. Woodv. 197.

Common Elm. April. Pen. Di. Hedges.

MARCH.

montána.

U. Leaves doubly ferrated, unequal at the base, egg-shaped but taper-pointed; fmooth.

With. Arr. 279.

April.

Pen. Di.

Wych Hazel Broad-leaved Elm. Shady lanes, outskirts of woods, about Kendal.

SAMBU'CUS. Cal. 5-toothed: Bloff. regular, with 5 shallow clefts: Berry juicy, closed, many-feeded.

nigra.

Tufts with 5 divisions: leaves winged: leafits nearly egg-shaped, serrated: stem tree-like May. With. Arr. 316. Woodv. 78.

Bloff. white: Berries black.

Pen. Tri.

Common Elder. White's Nat. Cal. Mar. 13.

SCIL'LA Bloff. 6 petals, expanding, deciduous: Filaments threadshaped.

bifólia.

Root folid: flowers nearly upright, without floral leaves: root leaves 2, spear shaped.

With. Arr. 338. E. Bot. 24. Bloff. blue. Squill.

Hex. Mo.

VACCI'NIUM. Calyx fuperior: Bloff.

1 petal: Filaments fixed to the receptacle: Berry 4-celled, many-feeded: beneath, dimpled.

V. Bunches terminating, nodding: Vitis-idæ'us. leaves inverfely egg-shaped; edge very entire and rolled back; dotted underneath.

With. Arr. 371.

Bloff. pale pink.

Red Wortle Berries. April. Oct. Mo.

Mountainous fituations.

Po'Pulus. Flowers Male and Female in catkins on diffinct plants. Calyx scales ragged: Bloff. turban shaped, mouth entire, flanting.

Female fummit 4-cleft: Capf. 2-celled: Seeds many, downy.

P. Leaves nearly triangular, tooth- al'ba. ed, and angular: cottony underneath.

With. Arr. 375.

White Poplar. Abele Tree. Oct. Mo. Hedges, woods, and near brooks.

MARCH.

P. Leaves circular, toothed and trémula. angular, fmooth on both fides.

With. Arr. 375.

Asp. Aspen Tree. Moift woods, and in boggy ground.

P. Leaves rowel-shaped, tapering nigra. to a point, ferrated, fmooth, on both fides.

With. Arr. 376.

Oct. Mo. Black Poplar. Near rivers, and wet shady places.

DA'PHNE. See Gen. Ch. page 3.

D. Bunches of about 5 flowers, lauréola. axillary: leaves spear - shaped, fmooth. April. With. Arr. 377. E. Bot. 119.

Bloff. yellowish white.

Spurge Laurel, or Laurel Meze-Oct. Mo. reon. Woods and hedges.

> ADO'XA. Cal. cloven, beneath: Bloff. 4 or 5-cleft, superior: Berry 4-celled, united to the cup.

Oft. Mo.

A. With. Arr. 386. Curt. 2. lina. Ina.

Tuberous Moschatel. May.

Damp woods, and shady places. Oct. Tet.

CHRYSOPLE'NIUM. Cal. coloured, 4 or 5-cleft: Bloff. none: Capf. 2-beaked, 1-celled, many-feeded, partly beneath; division of the valves extending but half-way down.

C. Leaves alternate.

With. Arr. 401. E. Bot. 54.

Flowers bright yellow.

Alternate-leaved Sengreen. Wet Decan. Di. fhady woods.

April.

C. Leaves opposite.

With. Arr. 402. Curt. 2. opositifó-

Flowers bright yellow.

Golden Saxifrage. Opposite-leaved
Sengreen. May. Decan. Di.
Moist shady places.

STELLA'RIA. Cal. 5 leaves, expanding: Petals 5, mostly divided down to the base: Caps. 1-celled: Seeds many.

MARCH:

média. S. Petals deeply divided: leaves egg-heart-shaped: stems with a hairy ridge on the side.

With. Arr. 418. Curt. 1.

Decan. Tri. Common Chickweed. Alsine Média. Very common. Oct.

Holóstea. S. Leaves spear-shaped, very finely ferrated: petals cloven.

With. Arr. 417. Curt. 2.

Flowers white.

Decan. Tri. Greater Stichwort. May. Hedges, and woods.

Ox'ALIS. Calyx 5 leaves: Petals connected by the claws: Capf. 5-fided, opening at the corners.

Acetofel'la. O. Stalk with one flower: leaves 3 together: leafits inversely heart-fhaped, hairy.

With Arr. 430. Curt. 2. Woodv.

20.

Decan. Pen.

Petals white, veined with purple.

Wood Sorrel. Cuckow-bread.
Woods, shady hedges. Apr

Fa'Gus. Male and Female flowers on the same plant: Bloff. none.

Male. Calyx 5-cleft, bell shaped.
Fem. Calyx with 4 teeth, changing into a prickly 4 valved, 2-feeded capsule.

F. Leaves egg-shaped; indistinct- sylvatica. ly serrated.

With. Arr. 444.

Catkins, globular. Bark fmooth, white. April.

Beech Tree. Common Beech. Dode. Di. Woods and hedges, in a calcareous foil.

PRU'NUS. Calyx 5-cleft; beneath: Petals 5: Drupa 1-celled, closed at the top: Nut with projecting feams.

P. Fruit - stalks folitary: leaves spinosa. spear-shaped, smooth: branches thorny.

With Arr. 457. Woodv. 84.

Black Thorn. Sloe Tree. April. Ico. Mo. Hedges.

Anemo'ne. Cal. generally none: Petals 5 to 10: Capf. many, with awns or tails formed by the stile. nemorófa. A. Seeds pointed: leafits fnipt:
. flem with 1 flower.

With. Arr. 499. Curt. 2. E. Bot. 355.

Flowers fometimes double, and fometimes entirely of a purplish red colour. April.

Poly. Poly. Wood Anemone. Woods, hedges, hollow-ways.

White's Nat. Cal. Mar. 17-Ap. 22.

CAL'THA. Cal. none: Petals 5: Nectarics none: Capf. feveral: many-feeded.

palústris. C.

With. Arr. 511. Curt. 1.

Petals yellow, from 5 to 7.

Poly. Poly. Marsh Marigold. Meadow-bouts.

Moist meadows, banks of rivers,
and pools. May.

The As'Pi. Pouch inverfely heartfhaped, notched at the end: valves like a keeled boat, often winged with a border: cells many-feeded. T. Pouches compressed, triangu- Burfa paslarly inversely heart - shaped, fmooth, without a border: rootleaves wing-cleft.

With. Arr. 571. Curt. 1.

Petals entire white. Sept.

Shepherd's Purfe, or Pouch. A- Tetd. Silicu. mong rubbish, road-sides.

DRA'BA. Pouch entire, ellipticaloblong, rather compressed: Valves flat, parallel to the partition: Style none.

D. Stalks naked: leaves sparingly verna. ferrated: petals divided.

With. Arr. 565. Curt. 1.

Common Whitlow grafs. Nail- Tetd. Silicu. wort. Walls, and dry places.

April.

LEPID'IUM. Pouch notched at the end, compressed: Valves sharply keeled; I feed in each cell.

Leaves winged, very entire: petræ um. petals notched, fmaller than the calyx.

With, Arr. 567. E. Bot. 111.

One of our fmallest, delicate plants.

Mountain Dittander. April.

Tetd. Silicu. Rocks, walls, and stony places.

CARDAM INE. Pod long, 2-edged, opening with a jerk: Valves rolling back parallel to the membranaceous partition: Summit a knob, entire: Calyx rather open.

hirsúta. C. Leaves winged, leafits opposite, stamens 4.

With. Arr. 578. Curt. 4. June.

Hairy-leaved Ladies Smock.

Tetd. Siliq. Gravelly foil, on the fides of rivulets.

Tussila'Go. See Gen. Ch. page 15.

Petasítes. T. Panicle egg-shaped: female florets few.

With. Arr. 720. Curt. 2.

Syng. Super. Butter-bur. Coltsfoot. April.

Moift meadows, pastures, banks
of rivers.

Bel'lis. Recept. naked, conical: Down none: Calyx hemispherical, with equal scales: Seeds inversely egg-shaped.

B. Stalk naked.

peren nis.

Sept.

With. Arr. 733. Curt. 1.

Common Daify.

Meadows and pastures.

Syng. Super.

APRIL.

CALLI'TRICHE. Cal. none: Petals 2: Seeds 4, naked, with a membranaceous border on one fide.

C. Upper leaves oval; not notched at the end.

With. Arr. 5. August.

Bloff. fmall, white.

Vernal Star-grafs. Water Fennel. Mo. Mo. Ditches, ponds, flow streams.

> OR'CHIS. Nectary like a horn, behind the flower: Bloff. gaping.

mas cula. Lip of the nectary 4-lobed, finely fcolloped: horn blunt: upper petals turned back. With. Arr. 24. Curt. 2. Woodv. 90.

Flowers purple.

May.

Early Orchis. Male Fool-stones. Meadows and pastures:

White's Nat. Cal. April 21.

Di. Mo.

VERO'NICA. See Gen. Ch. page 5.

V. Spike terminating: flowers fo-ver'nz. litary: leaves with finger-like divisions: fruit-stalks shorter than the calyx.

With. Arr. 13. E. Bot. 25.

Bloff. blue, green at the base.

Vernal Speedwell. May.

Dry barren foil, old walls and Di. Mo. rocks.

V. Flowers folitary: leaves with triphyllos. finger like divisions: fruit-stalks longer than the calyx.

With. Arr. 17. Curt. 6.

Bloff. blue. May.

Right or Upright Chickweed Tri- Di. Mo. ... fid Speedwell. Sandy fields.

SA'LIX. Catkin, each scale containing 1 flower: Bloff. none.

Male, a nectariferous gland at the bottom of the flower.

Fem. Style cloven: Capf. 1 cell, 2 valves: Seeds downy.

monan dra. S. Leaves strap-spear shaped: upper leaves oblique: catkins downy: flowers with 1 stamen. Curt. 6. With, Arr. 45. May.

Di. Mo. Rose Willow. Purple Willow. Hedges, watery places.

flowers with 3 stamens, sometimes 2.

With. Arr. 45.

Di. Mo. Smooth Willow.

Woods, hedges, banks of rivers.

pentan'dra. S. Leaves egg-shaped, acute: flowers with 5 stamens.

With. Arr. 46.

Di. Mo. Sweet Willow. Bay-leaved Willow. Woods and hedges.

amygdalinía. S. Leaves spear-shaped on leafstalks: stipulæ in shape of an irregular square.

With. Arr. 47. August.

Di. Mo. Almond-leaved Willow.

Willow beds, banks of rivers.

S. Leaves egg-spear-shaped: leaf- frágilis, stalks toothed with glands.

With. Arr. 47. Woodv. 198.

Crack Willow. Woods, banks of Di. Mo. rivers, hedges.

S. Leaves strap-spear-shaped, a- rúbra. cute.

With. Arr. 49. May.

Red Willow. Willow-beds. Di. Mo.

S. Leaves egg-oblong, smooth a- répens. bove, filky underneath. May. With. Arr. 51. E. Bot. 183.

Creeping Willow. Wet fandy Di. Mo. heaths.

S. Leaves oblong-egg-shaped, some-cinérea. what woolly underneath: stipulæ half-heart-shaped.

With. Arr. 54. Di. Mo.

Sallow Willow. Hedges, in a moift foil.

S. Leaves spear-strap-shaped, very viminalis. long, acute, filky underneath: branches rod-like.

With Arr. 54. May. Di. Mo.

Ofier. Willow-beds, woods, hedges.

al ba.

S. Leaves spear-shaped, tapering to a point, serrated, downy on both sides: the lowermost serratures glandular.

With Arr. 55.

Di. Mo.

White, or Common Willow. Woods, hedge rows, and wet meadows.

O'rhrys. Nectury flightly keeled underneath.

araniféra.

O. Bulb roundish: stem leafy: lip of the nectary roundish, entire, nicked, convex: longer than the petals.

With. Arr. 39. Curt. 6. E. Bot. 65.

Di. Mo.

Spider Ophrys. Chalky meadows, and pastures. May.

VALERIA'NA. Cal. none: Bloff. 1 petal, superior, bulging on one fide at the base: Seed 1, or else a 3 celled capsule.

locústa.

V. Stem forked: leaves strap-shap-ed, very entire.

With. Arr. 66. Curt. 5.

Bloff. bluish white. Leaves and

Stems fringed at the edges with fine white hairs.

Var. 4. Upper leaves ferrated.

July.

Lamb's Lettuce. Corn Sallad. Tri. Mo. Lettuce Valerian. Corn fields.

Holosteum. Calyx 5 leaves: Petals 5: Capf. 1-celled, nearly cylindrical, opening at the top.

H. Flowers in umbels.

umbellátum.

With. Arr. 175. E. Bot. 27.

Whole plant smooth. Fruit-stalks terminating mostly hanging down, of various lengths, each with 1 flower.

Petals pale red. May.

Umbelliferous Moufe Ear. Old Tri. Tri. walls, banks, fandy corn-fields.

EM PETRUM. Male and Fem. flowers on separate plants. Calyx with 3 divisions: Bloff. 3 petals. Male Stam. long. Fem. Berry with 9 seeds.

E. Stems trailing. May. nigrum. With. Arr. 176.

Tri. Enne.

Black-berried Heath. Black Crowberries. Crake-berries. Moist mountains, high heaths.

BETULA. See Gen. Ch. page 11.

al ba.

B. Leaves triangular-fpear-shaped, acute; smooth, doubly serrated.

With. Arr. 206.

Tet. Di. Bloff. egg. shaped, concave, green.

Birch Tree. Birk. May.

Woods, moist hedges.

Bux'us. Male and Fem. flowers on the fame, or on a different plant.

Male. Cal. 3-leaved: Bloff. 3 petals: Capf. 3-celled: 3-beaked: Seeds 2.

Fem. Cal. 4-leaved: Bloff. 3 petals: Capf. 3-celled, 3-beaked: Seeds: 2.

femper-virens. B. Male and Fem. flowers on the fame plant.

With. Arr. 210.

Leaves oval, thick, gloffy. Bloff. greenish white.

Common Box.

Tet. Tri.

Var. 2. Narrow-leaved.
On box-hills, woods, hedges.

SAGINA. Cal. 4-leaved: Petals 4: Capf. 1-celled, 4-valved, manyfeeded.

S. Stem upright, generally supericta. porting a fingle flower. May.

With. Arr. 216. Curt. 2.

Leaves strap-spear-shaped. Bloff. white.

Least Stichwort. Upright Pearl- Tet. Tet. wort. Gravelly foil.

LITHOSPER'MUM. Bloff. funnelfhaped, tube long, flender, open, and without valves at the mouth: Cal. with 5 divisions: Nuts 4, very hard, imperforated.

L. Seeds fmooth: bloffom much longer than the cup.

With. Arr. 226. E. Bot. 117.

purpúro ' cærúleum.

Bloff. purple. May.

Creeping Groomwell. Mountain Pen. Mo. ous and woody pastures.

Pen. Mo.

APRIL.

ASPERU'GO. Cal. when in fruit, the fides flat, parallel, indented.

procumbens. A. Calyx when in fruit compressed.

With. Arr. 231.

Bloff. fmall, blue. May.

Trailing Catchweed. Roads, and amongst rubbish.

PRI'MULA. See Gen. Ch. page 1.

P. Leaves wrinkled and toothed:
falk many-flowered: outermost
flowers drooping: border of the
blossom flat.

With, Arr. 234.

Bloff. pale yellow: leaf-stalks shorter than the leaf. May.

Pen. Mo. O. lips. Great Cowflips. Oxlip

Primrofe. Woods, hedges;
fometimes pastures.

officinális. P. Leaves wrinkled and toothed:
ftalk many-flowered: all the
flowers drooping: border of the
bloffom concave.

With. Arr. 234. Curt. 6. E Bot. 5.

Bloff. fweet-scented, full yellow, with an orange blotch at the base of each segment.

Cowslips. Cowslip Primrose. P Meadows and pastures, in a loamy clayey soil. May.

Pen. Mo.

CY'CLAMEN. Bloff. wheel-shaped, reflected: tube very short; mouth projecting: Recept. sleshy, supporting a berry inclosed in a case.

C. Bloffom bent back as if broken: europæ'um. leaves circular, fcolloped.

With. Arr. 235.

Leaves greatly varying from Pen. Mo. round to angular. Bloffom purplish pink.

RHA'MNUS. Cal. tubular: Scales projecting the stamens: Bloff. none: fruit a drupa.

R. Thorns terminating: flowers cathárticus.

4-cleft, male and female on different plants: leaves egg-shaped:
stem upright.

With, Arr. 256. Woodv. 114.

Pen. Mo.

APRIL.

Pen. Mo. Purging Buckthorn. May.

Woods and hedges near Norwich.

Fran'gula. R. Without thorns: flower hermaphrodite, with 1 pistil: leaves very entire.

With. Arr. 259. E. Bot. 250.

Berry with 3 cells: 2 feeds.

Black berry-bearing Alder. Alder Buckthorn. May.

Woods, wet hedges.

RI'BES. Petals 5, they and the stamens fixed to the calyx: Style cloven: Berry beneath, 1-celled, many-seeded.

fpicatum. R. Spikes upright: petals oblong:
floral-leaves shorter than the
flowers.

With. Arr. 265.

Pen. Mo. Its floral leaves being shorter than the flowers, will prevent its being taken for the R. alpinum. May.

R. Branches prickly: fringe of the Groffulária. · leaf-stalks hairy: berries hairy. With. Arr. 266.

Rough Goofeberry. Fenberry. Hedges.

Pen. Mo.

Branches prickly: berries Uva-crifpa. fmooth: fruit-ftalks with a flower-scale of 1 leaf.

With. Arr. 266.

A triple thorn beneath the buds. May. Pen. Mo. Smooth Goofeberry. Woods, hedges.

- FRITILLA RIA. Bloff. with 6 petals or 6 divisions, bell-shaped: Nectary a cavity above each claw: Stam. the length of the bloffom: Summit 3-cleft: Capf. Superior, 3-celled.
- F. All the leaves alternate: stem Meléagris. 1-flowered.

With. Arr. 335. Curt. 3.

Fritillary. Chequered Daffodil. Meadows, pastures.

May. Hex. Mo.

APRIL.

- fylvístris. Tu'lipa. Bloff. 6 petals: bellshaped: Style none: Capf. superior, 3-celled.
 - T. Flower rather drooping: leaves fpear-shaped: stamens hairy above the base.
- Hex. Mo. With. Arr. 336. E. Bot. 63.

 Bloff. yellow sweet-scented.
 Old chalk-pits in Suffolk and
 Norfolk. Tulip.
 - ORNITHO GALUM. Bloff. 6 petals, upright, permanent, above the middle expanding: Filaments, the alternate ones dilated at the base: Caps. superior, 3-celled.
- lúteum. O. Stalk angular, 2-leaved: fruitftalks in an unbranched umbel. With. Arr. 336. E. Bot. 21.
- Hex. Mo. Yellow Bethlem Star. Moist fandy places.
- umbellátum. O. Flowers forming a corymbus: outer fruit-stalks taller than the central ones.

With. Arr. 337. E. Bot. 130.

Hex. Mo. Common Bethlem Star. Woods, and meadows.

HYACIN'THUS. Bloff. bell-shaped, permanent; fegments rolled back: Germen with 3 nectariferous pores at the top.

H. Bloff. tubular bell-shaped, with nonscriptus. 6 divisions, segments rolled back: floral leaves in pairs.

With. Arr 343. Curt. 18. E. Bot. 377.

Flowers varying from blue, to white or flesh colour.

Harebell. English Hyacinth. Woods and hedges. May.

Hex. Mo.

SCILLA nutans. See E. Bot. 377.

VACCINIUM. Calyx Superior: Bloff. 1 petal: Filaments fixed to the receptacle: Berry 4-celled, manyfeeded; beneath, dimpled.

V. Fruit-flalks 1-flowered: leaves Myrtillus. ferrated, egg-shaped: stem regular. With. Arr. 370.

May.

Bloff: nearly globular, a little flatted at the base: pale reddish purple.

Black Worts. Black Whortle-ber- Oct. Mo. rics. Hurtle berries. Bilberries. Woods and heaths.

uliginósum. V. Fruit-stalks 1-flowered: leaves very entire, inversely egg-shaped, blunt, smooth.

With. Arr. 370.

Bloff. pink.

May.

OA. Mo.

Great Bilberry-bush or Whortle. Moistish woods, heaths, and pastures.

Oxycoc'cos. V. Fruit-stalks single, or in pairs: leaves egg-shaped, very entire; edges rolled back: stem threadshaped, trailing, not hairy.

With. Arr. 372. E. Bot. 319.

Bloff. deep flesh-colour.

OA. Mo. Cran-berries. Moss-berries. Moorberries. Fen-berries.

Peaty bogs, in the north.

A'CER. Male flowers intermixed:

Calyx 5-cleft: Bloff. 5 petals:

Capf. 2 or 3; 1-feeded: ending in a leaf-like expansion.

Pseudo-plá- A. Leaves 5-lobed, blunt, unequally serrated: flowers in compound, pendent bunches.

With. Arr. 368. E. Bot. 303.

Bloff. yellow green.

Sycamore Tree. Sycamore Maple. Oct. Mo. Woods, hedges. June.

Poly'gonum. Cal. none: Bloff. refembling a cup with 5 divisions: Seed 1, angular; generally naked.

P. Flowers axillary: leaves spear-aviculare. shaped: stem trailing.

With. Arr. 383. Curt. 1.

Bloff. greenish on the outside, white within. Sept. Oct. Tri. Knot-grass. Snakeweed.

QUER'CUS. Male and Fem. flowers on the fame plant: Bloff. none.

Male. Calyx 5-cleft: Stamens from 5 to 10.

Fem. Cal. 1 leaf, very entire, rough: Styles from 2 to 5:
Nut egg-shaped, leather-like, of 1 seed, and when ripe of 1 cell.

Q. Leaves on leaf-stalks, oblong, Róbur. broadest towards the end, indent-

ations rather acute, angles blunt: fruit nearly fitting.

With. Arr. 387. Woodv. 126.

Oct. Tet. Bay Oak. About Newbury.

fémina. Q. Leaves oblong, on short leafflalks, blunt wing-clest, with indentations: fruit mostly solitary, on long fruit-stalks.

Oct. Tet. With. Arr. 387. Woods and hedges.

MERCURIA'LIS. Male and Female flowers on different plants. Bloff. none: Cal. with 3 divisions.

Male Stamens from 9 to 12. Anthers globular, double.

Fem. Capf. double, 2-celled, 1 feed in each.

peren'nis. M. Stem undivided: leaves rough.

Flowers yellow green.

Enne. Di. Dog's Mercury. Woods, hedgebanks. With. Arr. 392. Curt. 2. May.

SAXI FRAGA. Cal. 5-toothed: Bloff. 5 petals: Capf. beneath 2-beak-

ed; 2 celled, many-feeded, opening with a hole between the ftyles.

S. Leaves egg-shaped, scolloped, nivális. nearly sitting: stem leastless: flowers in heads.

With. Arr. 403. Oct.

Mountain Saxifrage. Summits of Dec. Di. high mountains.

S. Stem-leaves opposite, tiled; the oppositistem-leaves opposite, tiled; the opposite opp

With Arr. 404. Curt 6. E. Bot. 9. Petals very blunt, bright purple. Heath-like Sanifrage. Rocks.

June.

S. Stem-leaves kidney - shaped, granuláta. gashed: stem branched: root beaded.

With. Arr. 405. Curt. 1. Woodv. 232.

Bloffoms white. May. Dec. Di.

White Sa ifrage. Dry meadows, pastures.

S. Stem-leaves wedge-shaped, al- tridactylites, ternate, 3-cleft: stem branched:

maritima.

B

leaves in very dry fituations, fometimes undivided.

With. Arr. 406. Curt. 2.

Turns red after flowering.

Dee. Di. Rue Whitlow-grass. Rue-leaved Saxifrage.

fhaped, with 3 divisions: fegments mostly 3-cleft: stem limber, lying down, much branched: fruit-stalks very long, 1flowered.

With. Arr. 890.

Dec. Di. Flower yellow white, with a greenish tinge. June. Rocks.

SILE'NE. Cal. diftended: Petals 5, with claws: crowned at the mouth: Capf. 1 or 3-celled.

S. Stem trailing, branches ascending: Calyx veined, smooth: Petals cloven, segments very entire.

With. Arr. 414.

Petals white.

August.

Sea Campion, or Catch fly. Sandy Dec. Tri. places on the fea-coaft.

CERA'STIUM. See Gen. Ch. page 12.

C. Upright, woolly and clammy. viscosum. With. Arr. 433. Curt. 2.

Petals white, shorter than the calyx.

Clammy Moufe-Ear. Oct.

Meadows, walls, dry banks, ant- Dec. Pen. hills.

C. Flowers with 5 flamens: petals femidenotched at the end.

With. Arr. 435. Curt. 2.

Flowers white.

Dec. Pen.

Leaft Mouse-Ear. May. Walls, pastures, heaths.

PRU'NUS. See Gen. Ch. page 25.

P. Umbels mostly on short fruit- Cérasus. stalks: leaves egg-spear-shaped, smooth, doubled together.

With. Arr. 455.

E

Bloff. white. Fruit red.
Common Wild Cherry-Tree.
Woods, hedges. May.

doméftica. P. Fruit-stalks mostly solitary:
leaves spear-egg-shaped, coiled:
branches thornless.
With. Arr. 456. Woodv. 85.

Ico. Mo. Common Plumb-Tree. Hedges.

egg-shaped, slightly woolly, coiled: branches with thorns.

Ico. Mo. With. Arr. 456.

Bullace Plumb. Hedges.

CRATE GUS. Cal. 5-cleft: Petals 5: Berry beneath, opening at the top, 1-celled, 1 or more feeded.

Oxyacantha. C. Leaves mostly 3-cleft: fegments blunt, ferrated.

With. Arr. 459.

Ico. Di. White-thorn. Hedges, woods, old parks. June. White's Nat. Cal. Ap.18—May 12.

Son' BUS. Cal. 5-cleft: Petals 5: Pomum, 5-celled, open at the top, 3-feeded. S. Leaves winged, finooth on both aucupária. fides.

With. Arr. 460. E. Bot. 337.

Flowers whitish.

Quicken Tree. Mountain Ash.

Woods, hedges, in mountainous Ico. Tri. and boggy fituations.

S. Leaves winged, woolly under- doméstica. neath.

With. Arr. 460. E. Bot. 350.

True Service, Sorb. Mountains. Ico. Tri.

Py'Rus domestica. See E. Bot. 350.

Py'rus. Cal. 5-cleft: Petals 5: Pomum, beneath: 5-celled; many feeded.

P. Leaves ferrated, fmooth: flow-commúnis. ers forming a corymbus.

With. Arr. 462.

Bloff. white. May. Ico. Pen. Pear Tree. Woods, hedges.

POTENTIL'LA. Cal. 10-cleft: Petals 5: Seeds roundish, naked, wrinkled, fixed to a receptacle, which is small, juiceless, spongy, tubercled. ver na.

P. Root-leaves 5 together, sharply ferrated, dented at the end: Rem-leaves 3 together: stem declining.

With. Arr. 475. E. Bot. 37.

Petals yellow, nicked, with fometimes a tawny fpot at the base.

Ico. Foly.

- Spring Cinquefoil. Rocky mountainous pastures and barren meadows.
- POTE RIUM. Stamens and Piftils in different flowers, on the same plant: Cal. 4 leaves: Bloff. with 4 divisions.

Male. Stam. 30 to 40.

Fem. Drupa juiceless; beneath; 1 or 2-celled; formed of the indurated tube of the bloffom.

Sanguifórba.

P. Thornless: stems somewhat angular.

With. Arr. 493. Curt. 2.

Flowers greenish, sometimes purplish on the outside.

Upland Burnet. Poly. Di.

May.

In a dry calcareous foil.

ANEMO'NE. See Gen. Ch. page 25.

A. Leaves doubly winged; petals Pulfatil'Ia. ftraight.

With. Arr. 498. E. Bot. 51.

Petals deep purple. May.

Pasque Flower. High pastures. Poly. Poly.

A. Leaves doubly winged: petals praten fis. the end turned back.

Flower small, dark purple.

With. Arr. 498. Woodv. 148.

Dark-flowered Anemone. Dry Poly. Poly. hilly pastures.

A. Seeds pointed: leafits fnipt: apennina. petals fpear-shaped, numerous.

With. Arr. 500. Curt. 6.

Petals blue: fometimes elliptical.

Mountain Anemone. Woods and Poly. Poly. flady places.

RANUN'CULUS. See Gen. Ch. p. 7.

R. Root-leaves kidney-shaped, scolloped, cut: stem-leaves singered, strap-shaped: stem many flowered, With. Arr. 505. Curt. 2. Bloff. yellow.

Poly. Poly. Sweet Wood Crowfoot. Goldilocks. Woods, hedges, in a clayey foil.

A juga. Bloff. upper-lip very small: Stamens longer than the upper-lip.

Chamæ'pithys.

A. Leaves 4-cleft, ftrap-shaped;
very entire: flowers fitting, lateral, folitary: stem spreading.

With. Arr. 516. E. Bot. 77.

Bloff. yellow, with minute red dots. June.

Didy. Gym. Ground Pine. Germander. Sandy fallow fields.

GLECO'MA. Cal. 5-cleft: Anthers in pairs, each pair forming a cross.

Hederácea. G. Leaves kidney-shaped, scolloped. With. Arr. 526. Curt. 2. Woodv. 28.

Bloff. blue. May.

Didy. Gym. Ground Ivy. Gill. Cat's-foot. Groves, hedges, shady places.

LAMIUM. See Gen. Ch. page 8.

L. Leaves deeply and irregularly diffectum, cut; stem-leaves extending down the leaf-stalks.

With. Arr. 527.

Dead-nettle. Kitchen gardens, Didy. Gym. fallow-fields, bank between Pimlico and Chelsea. Sept.

LATHRÆ'A. Cal. with 4 or 5 divisions: Germen with a depressed gland at the base of the seam: Caps. 1-celled: Receptacles lateral, sponge-like.

L. Stem undivided: flowers pen- Squamária.
dant: lower lip 3 cleft.

With. Arr. 547. E. Bot. 50.

Bloff. pale purple, except the lower lip.

Great Toothwort. Woods, shady Didy. Ang. lanes, bottom of mountains.

May.

Pedicula Ris. Calyx 5-cleft: Capf. 2-celled, sharp-pointed, slanting: Seeds few, coated.

P. Stem branched: calyx oblong, sylvatica.

angular, fmooth: lip of the bloffom heart-shaped.

With. Arr. 548.

Bloff. purple.

Didy. Ang. Common Loufewort.

Wet pastures and heaths. July.

SCROPHULA'RIA. Cal. 5 - cleft:

Bloff. with 5 divisions: tube globular: the lower fegment reflected: Caps. 2-celled, partition
double.

vernális.

S. Leaves heart-shaped; those of the stem in threes; fruit-stalk axillary, solitary, cloven.

With. Arr. 554.

Bloff. egg-shaped yellow.

Didy. Ang.

Yellow Figwort. May. Watery places, hedges.

THLA'SPI. See Gen. Ch. p. 26.

perfoliátam. T. Pouches inverfely heart-shaped, frem - leaves heart - shaped, frmooth, somewhat toothed: petals as long as the calyx: stem branched.

With. Arr. 569.

Bloff. white.

Perfoliate Shepherd's Purfe. Stone- Tetd. Silicu. pits.

COCHLEA'RIA. Pouch notched at the end, turgid, rough, manyfeeded; valves bulging, blunt.

C. Root-leaves heart-circular; officinális. flem-leaves oblong, a little indented.

With. Arr. 571. Woodv. 29.

Petals fleshy clear white.

Scurvy Grass. May. Tetd.Silicu. Sea shores, inland mountains.

DENTA'RIA. Pod cylindrical, bordering, opening with a jerk: valves rolling back, rather shorter than the sponge-like partition: Summit notched: Cal. close.

D. Lower leaves winged: the up-bulbifera. per ones undivided.

With. Arr. 576. E. Bot. 309.

Flowers flesh-coloured.

Bulbed Coralwort. Woods, and Tetd. Silique shady places. May.

CARDAM INE. See Gen.Ch. page 28.

amára.

C. Leaves winged: fuckers from the bosom of the leaves: leafits of the stem-leaves angular, sitting.

With. Arr. 580. Curt. 3.

Flowers white.

May.

Tetd. Siliq.

Bitter Crefs, or Ladies Smock. Near purls of water, rivulets.

praten fis.

C. Leafits of the root-leaves roundish, those of the stem-leaves spearshaped: very entire.
With. Arr. 579. Curt. 3. Woodv.
30. May.

Flowers purplish.

Tetd. Siliq.

Common Ladies Smock. Meadows, moist pastures.

ERY'SIMUM. Pod straight, strap-shaped, exactly 4-sided: Cal. closed: Summit a knob.

Barbárea.

E. Pods indistinctly 4-cornered; Leaves lyre-shaped, the terminating segment circular.

With, Arr. 584.

Bloff. yellow.

Oa.

Winter Creffes, or Rocket. Walls, Tetd. Siliq. banks of running streams.

E. Leaves heart-shaped. With. Arr. 584. Curt. 2. Woodv.

Alliária.

245.

Flowers in a corymbus.

Jack by the Hedge. Sauce alone.

May. Tetd. Siliq.

Hedges, ditch banks.

GERA'NIUM. Cal. with 5 divisions:

Bloff. 5 petals: Pift. 1: Summits
5, common receptacle beaked:

Seeds 5, with a very long twisted tail.

G. Fruit-stalks and floral-leaves mol'le. alternate: petals cloven: calyx awnless: stem somewhat upright.

With, Arr. 604. Curt. 2.

Flowers purplish red. Oct. Mond. Dec. Soft Crane's-bill. Dove's-foot. Dry

banks.

G. Leafits by fives or by threes: robertiánum.

APRIL.

lobes wing-cleft: calyx with 10 angles.

With. Arr. 608. Curt. 1.

Bloff. red, sometimes white.

Mond. Pec. Herb Robert. Stockbill. August. Walls, hedges, rubbish, stony places.

cicutárium. G. Flowers in umbels: leafits fitting, wing cleft, blunt: petals entire.

With. Arr. 609. Curt. 1. Flowers rose red. Oct.

Mond. Dec. Hemlock-leaved Crane's-bill.

Corn-fields, road-fides, in a fandy foil.

Fuma'ria. Cal. 2-leaved: Bloff. gaping: Filaments 2, membranaceous, each supporting three anthers: Caps. 1-celled; manyfeeded.

intermédia. F. Root bulbous folid: stem simple: sloral leaves hand-shaped, as long as the blossom.

With. Arr. 620.

Flowers purplish red, and yellow.

Fumitory. May. Diad. Hex. Levan's Park, near Kendal.

F. Legumens strap-shaped, 4-sided; capnoides. stems spreading, the angles acute.

With. Arr. 620.

Flowers whitish. Old walls, near Diad. Hex. Castleton, Derbyshire.

O'ROBUS. Cal. blunt at the base: the 2 upper-teeth shorter, but more deeply divided: Style threadshaped.

O. Leaves winged, spear-shaped, tuberòsus. stipulæ half-arrow-shaped, very entire: stem simple.

With. Arr. 631. Curt. 1.
Bloff. purple, 3 or 4 together.

Heath Peafeling. May. Diad. Dec. Moist heaths, woody meadows.

Vi'cia. Summit bearded across underneath: Legumen generally with knot-like protuberances.

V. Legumens mostly in pairs, up-fativ'a. right: leasits inversely spear-shaped, blunt, notched: stipulæ toothed: seeds compressed.

With. Arr. 636. E. Bot. 334.

Bloff. reddish purple.

Diad. Dec. Common Vetch. Tare. May.

Lanes in the valleys of Dartmoor.

Tussila'Go. See Gen. Ch. page 15.

hybrida. T. Panicle oblong: female flowers numerous without bloffoms.

With. Arr. 721.

Syng. Super. Long-stalked Colt's - foot. Moist meadows, banks of the river Irwell.

MAY.

A'PHANES. Cal. tubular: rim flat, 4 or 8-cleft: Bloff. none: Seed 1; egg-shaped, but tapering to a point.

A. Leaves 3-lobed; lobes with 2 arven'fis. or 3 clefts.

With. Arr. 4.

Flowers axillary; fmall, greenish white.

Parsleypiert. Corn-fields, and dry Mo. Mo. gravelly lands. August.

HIPPU'RIS. Cal. none: Summit fimple: Seed 1.

H. Leaves awl-shaped: 8 in a vulgáris. whirl.

With. Arr. 5. Curt. 4.

Common Mare's-tail. Paddow- Mo. Mo. pipe. Muddy ponds and ditches.

May.

VERO'NICA. See Gen. Ch. page 5.

fhaped, fitting: floral leaves fpearfhaped; longer than the flowers. With. Arr. 12. Curt. 2.

Bloff. pale blue.

Di. Mo. Wall Speedwell. Chickweed. Old walls, rubbish, fallow-fields.

officinális. V. Spikes on lateral fruit-stalks: leaves opposite: stem trailing.

With. Arr. 13. Curt. 3. Woodw.

219.

Bloff. purplish blue. Aug:

Di. Mo. Common Speedwell. Male speedwell. Fluellin. Barren ground, heaths.

alpina. V. Corymbus terminating: leaves opposite: cups rough with hair.

With. Arr. 14.

Di. Mo. Bloff. fmall, not a fine blue.

Mountains.

ferpyllifólia. V. Bunch terminating, fomewhat fpike-like: leaves egg-fhaped, fmooth, fcolloped, 3-fibred: ftamens hardly longer than the bloffom.

With. Arr. 14. Curt. 1.

Bloff. pale blue: upper fegment with 5 or 8 purple lines, lateral fegments with 3: smallest fegment white.

Smooth Speedwell. Paul's Betony. Di. Mo. Meadows and pastures.

V. Bunch terminating: leaves humifusa. heart-shaped but roundish, slightly scolloped: stems creeping.

With Arr. 15. Di. Mo.

High mountains, wet shading rocks.

V. Bunches lateral, of few flow-montána. ers: cup rough with hair: leaves wrinkled, egg-shaped, scolloped, on leaf-stalks: stem feeble. With. Arr. 16. Curt. 4.

Bloff. purplish, streaked. June. Di. Mo. Mountain Madwort. Stalked Speedwell.

V. Bunches lateral: leaves egg- Chamædrys shaped, sitting, wrinkled, toothed: stem with 2 opposite rows of hairs.

With. Arr. 16. Curt. 1.

it

1,

ne

1.

Bunches frequently opposite.

Bloff. a fine blue.

Di. Mo. Wild Germander. Germander Speedwell.

Pastures, fides of hedges.

PINGUI CULA. Bloff. gaping, ending in a spur: Cal. 2-lipped, 5cleft: Capf. 1 cell.

vulgáris. Nectary cylindrical, as long as the petal.

With. Arr. 18. E. Bot. 70.

Bloff. violet, purple, and reddish, with white lips, and an ashcoloured woolly fpot on the palate.

Di. Mo. York fhire Common Butterwort. Sanicle. Bogs.

OR'CHIS. See Gen. Ch. page 30.

O. Lip of the nectary 4-cleft, finely fcolloped: horn blunt, afcending: petals blunt, approaching. Lip of the nectary with 3 lobes, the middlemost notched at the end.

With. Arr. 23. Curt. 3. June.

Mório.

Flowers purple, rather few. The green lines on the petals diftinguish it from every other Orchis.

Var. 1. Flowers flesh colour, or Di. Mo. white.

Meadow Orchis. Moist meadows.

O. Lip of the nectary with 4-clefts usuláta. rough with dots: horn blunt: petals distinct.

With. Arr. 25. E. Bot. 18.

Petals purplish without, greenish white within. Spike short, seldom an inch. Lip white, with beautiful purple dots, not rough, as mentioned by Linnæus.

Dwarf Orchis. June. Di. Mo. Dryish meadows and pastures.

O. Lip of the nectary 5-cleft: militáris. rough with dots: horn blunt: petals confluent.

Flowers in a spike, purplish or ash colour. Petals growing together. Spur short, bluntish.

e

e.

Di. Mo.

datifolia.

Var. 2. Fusca. Lip 3-cleft, fegment very broad, notched at the end. With a greenish cast.

With. Arr. 25. Curt. 6. E. Bot. 16.

Var. 3. Lip 3-cleft, fegments ftrap-shaped.

Man Orchis. June. Meadows in a chalky foil.

O. Bulbs fomewhat hand-shaped, straight: horn of the nectary co-nical: lip with 3 lobes; the lateral ones bent back: floral-leaves longer than the flowers.

Bulbs straight, not much straddling, divided into 2 or 3 singers. Leaves, especially the lower, a little spotted. Differs from O. maculata, as follows. Floral-leaves longer than the slowers. Leaves nearly without spots. Stem hollow.

Var. 2. Middle fegment of the lip blunt, as narrow again as the fide ones, but of the fame length.

Var. 3. Middle fegment of the Di. Mo. lip egg-shaped, pointed, as long again as the side ones.

Flowers purple, the throat whitish with purple dots.

With. Arr. 26. Curt. 5.

Broad - leaved, or Male - handed Orchis. Moist meadows, pastures, marshy grounds. June.

O. Bulbs thread-shaped: lip of abortivathe nectary egg-shaped: very entire: stem leastess.

With. Arr. 29.

Stem sheathed with scales. Flowers violet. Horn conical, slightly curved, very much pointed.

Purple Birds' - nest. Birds' - nest
Orchis June. Di. Mo.

Dryish meadows.

O'PHRYS. See Gen. Ch. p. 34.

O. Bulbs fibrous, bundled: ftem Nidus. ávisfheathed, leafleís: lip of the nectary cloven.

With. Arr. 32. E. Bot. 48.

ováta.

Di. Mo.

Whole plant brown. Stalk clothed with sheathing leaffcales.

Di. Mo. Birds'-nest Twayblade. June. Woods and shady places—rare.

O. Bulbs fibrous: stem with 2 leaves: leaves egg shaped: lip of the nectary cloven.

With. Arr. 34. Curt. 3.

Fowers with a fragrant, musky fcent.

Common Twayblade. Woods and moist shady thickets.

Var. 2. Leaves acute. Much fmaller and greener than the preceding.

Low wet grounds. June.

nusciféra. O. Bulbs roundish: stem leafy:
lip of the nectary convex, clothlike, with 3 divisions, the middle segment cloven.

With. Arr. 37. E. Bot. 64.

Flowers thinly scattered, 4 to 15, having much more the appearance of a fly than those of O. apifera of a bee. The blue

fpot upon the base of the middle segment of the nectary lip, contributes much to its resemblance to a fly.

Fly Orchis, or Twayblade. Meadows and pastures in a calcareous soil.

June. Di. Mo.

SERA'PIAS. Bloff. 6 petals, expanding: Nectary egg-shaped, bulging: Lip egg-shaped: Caps. beneath, 1-celled, 3-valved.

S. Leaves egg-spear-shaped, embracing the stem: flowers nearly horizontal: lip pointed, entire, rather shorter than the petals. With Arr. 40 E. Bot. 260.

Petals purplish green. Aug.

Common Helleborine. Bastard Di. Mo. Hellebore. Woods, groves, hedges.

LEM'NA. Male. Cal. 1 leaf: Bloff. none.

Fem. Cal. 1 leaf: Bloff none.

Style 1. Capf. 1 or 2-celled; 1 or 2-feeded.

MAY.

polyrhíza. L. Leaves fitting: root crowded. With. Arr. 44.

Di. Mo. Greater Duckmeat. Ditches. Sept.

Salix. See Gen. Ch. p. 31.

vitellina. S. Leaves egg-spear-shaped, acute: fmooth above; ferratures like griftle.

With Arr. 47.

Di. Mo. Yellow Willow. Ofier holts, frequent.

folia.

S. Trailing: leaves fpear-strapshaped, stiff and straight, nearly
sitting, cottony underneath.

With. Arr. 52.

Di. Mo. Rofemary-leaved Willow. Low fields.

reticulata. S. Leaves egg-shaped, blunt. With. Arr. 49.

Di. Mo. Network Willow. Mountains of Wales.

capréa. S. Leaves egg-shaped, wrinkled, cottony underneath, waved, tooth-Di. Mo. ed towards the end.

With. Arr. 53.

S. Leaves woolly on both fides, aurita. inverfely egg-shaped, furnished with appendages.

With. Arr. 50.

Round-eared Willow. Woods and Di. Mo. hedges. June.

S. Leaves egg-oblong, cottony underneath: upper ones entire, lower ones scolloped.

With. Arr. 53. Di. Mo.

Thickets and wet places.

S. Leaves woolly on both fides, lanáta. roundish, acute.

With, Arr. 50.

Valleys in the Highlands. Di. Mo.

S. Leaves egg-shaped, pubescent fus'ca. underneath.

With. Arr. 51.

Sand Willow. Brown Willow. Di. Mo. Sea shores amongst loose blowing fand.

Anthoxan'thum. Cal. husk of 2 valves, and 1 flower: Bloff. husk 2 valves, tapering to a point: Seed 1.

odorátum. A. Spike oblong-egg-shaped: florets longer than the awns, supported on short fruit-stalks.

With. Arr. 58. Curt. 1.

Bloff. double.

June.

Di. Di. Spring Grass. Sweet-scented.

Meadows and pastures.

VALERIA'NA. See Gen. Ch. p. 34.

rúbra. V. Flowers with a tail: leaves fpear-shaped, very entire.

With. Arr. 65.

Tri. Mo. Red Valerian. Old walls, rubbish. Aug.

dioica. V. Stameniferous and piftilliferous flowers on different plants: leaves winged, leafits very entire, fomewhat ferrated. Stameniferous plants with imperfect piftils.

With, Arr. 65. Curt. 4.

Tri. Mo. Small Valerian. June. Moist meadows.

BRYO'NIA. Cal. 5-toothed: Bloff. with 5 divisions.

Male. Anthers entire at the base.

Fem. Style 3-cleft: Berry roundith, mostly 1-seeded.

B. Leaves hand-shaped: rough on dioscaboth sides, with callous points: male and semale flowers on different plants.

With. Arr. 67. Woodv. 189.

Flowers yellow white. Berry red.

Red-berried Bryony. Wild Vine. Tri. Mo. Hedges, thickets.

ERIO'PHORUM. See Gen. Ch. p. 10.

E. Straws cylindrical: leaves flat: polystáchion fpikes on fruit-stalks.

With. Arr. 72. Curt. 4.

Cotton Grafs. June. Tri. Mo. Bogs, marshy grounds.

Mon'tia. Cal. 2 leaves: Bloff. 1 petal, irregular: Capf. 1-celled, 2-valved.

2-valved.

M. fontána.

With. Arr. 175. Curt. 3.

Bloff. white. Seeds black, shining.

Small Water Chickweed. Purstane. Tri. Tri.

Springs, watery lanes.

TILLEA. Cal. with 3 or 4 divifions: Petals 3 or 4 equal: Caps. 3 or 4-seeded.

muscósa. T. Trailing: flowers 3-cleft.

With. Arr. 175. E. Bot. 116.

Stems at first nearly upright, generally red.

Tri. Tri. Moffy Red-shanks. June Dry heaths. Norfolk, Suffolk.

Polycar'pon. Capf. 5 leaves: Petals 5, small egg-shaped: Capf. 1-celled; 3-valved.

lum. P. Stem branched, trailing: leaves in fours, inversely egg-shaped.

With. Arr. 176.

Sometimes without petals.

Four-leaved Allfeed. Aug.

Dry fandy ground.

SHERAR DIA. Bloff. 1 petal, funnelfhaped, long: Seeds 2, naked, crowned by the calyx.

ers terminating.

With. Arr. 185. Curt. 5.

Bloff. blue or purple.

Little Field Madder. Sept. Tet. Mo.

Corn, and fallow fields.

ASPERULA. Bloff. 1 petal, funnelfhaped: Seeds 2, globular, like cork.

A. Leaves 8 in a whirl, fpear-odoráta. fhaped: flowers in bundles, on fruit-stalks.

With. Arr. 185. Curt. 4.

Flowers a beautiful fnowy white.

Sweet Woodroof. Woods, shady Tet. Mo. places.

GALIUM. Bloff. 1 petal, bell-shaped, short: Seeds 2, nearly globular, beneath like cork.

G. Stem hairy, only branched at cruciátum, the base: leaves hairy, egg-shaped, 4 in a whirl: fruit-stalks with 2 or 3 leaves.

With. Arr. 186. E. Bot. 143.

Bloff. yellow. July.

Croffwort. Mugwort. Hedge- Tet. Mo. banks, rough places.

Aparine. G. Leaves 8 in a whirl, fpearfhaped: keel rough, with prickles pointing backwards: joints woolly.

With. Arr. 192. Curt. 2. Woodv. 269.

Tet. Mo. Bloff. white. June.

Catchweed. Goofegras. Cleavers.

PARIETA'RIA. Fem. florets mixed with hermaphrodite on the same branch: Cal. 4-cleft: Bloff: none: Seeds 1; superior lengthening.

officinális. P. Leaves spear-egg-shaped: fruitstalks forked: cup of 2 leaves. With Arr. 201. Curt. 4. Woodv. 142.

Stems reddish. Bloff. greenish white.

Tet. Mo. Common Pellitory of the Wall.

Old walls, rubbish. Sept.

Vis cum. Male and Fem. flowers on different plants: Bloff. none.

Male. Cal. with 4 divisions: Filam. none: Anthers fixed on the calyx.

Fem. Cal. 4 leaves, superior:

Style none: Berry pulpy; 1-celled, 1-fecded: Seeds heart-shaped.

V. Leaves spear-shaped, blunt: al'bum. stem forked: spikes axillary.

With. Arr. 203. Woodv. 270.

Bloff. greenish white.

Miffeltoe. Miffel. On apple-trees, Tet. Mo. pear, oak, hazel, &c.

BUFFO'NIA. Cal. 4 leaved: Bloff. 4 petals: Capf. 1 cell; 2 feeds.

B. Var. 2. tenuifolia. With. Arr. 205.

Flowers at the base of the leaves white.

Bastard Chickweed. Toadgrass. Tet. Di. Hounslow Heath, sea coast. June.

BETULA. See Gen. Ch. p. 11.

B. Leaves circular, scolloped. nána.

With. Arr. 207.

Mountains, wet heaths. Tet. Di.

Myrica. Flowers in catkins, on different plants. Cal. 2 leaves: Bloff. none.

Fem. Drupa 1-celled, fuperior: Seed 1.

Gále. M. Leaves spear-shaped: somewhat serrated: stem shrub-like. With. Arr. 208.

Tet. Di. Flowers appear before the leaves.

Sweet Gale, or Willow. Dutch

Myrtle.

I'LEX. Stam. and Pift. variously disposed: Cal. 4-toothed: Bloff. wheel-shaped: Style none: Berry 4-seeded.

Aquifólium. I. Leaves egg-shaped, acute, thorny, on leaf-stalks: flowers in a kind of umbel, axillary.

With. Arr. 210.

Flowers small, whitish. Common Holly.

Tet. Tet. Var. 2. Berries yellow. June. Woods, hedges, heaths.

POTAMOGETON. Cal. none: Petals 4: Style none: Seeds 4.

nátans. P. Leaves oblong egg-shaped, on leaf-stalks; floating.

Var. 1. floating.

With. Arr. 211.

Flowers fitting, numerous, yellowish.

Broad-leaved Pondweed. Ponds, Tet. Tet. flow rivers. Aug.

P. Leaves egg-shaped, tapering to den'sum.
a point: opposite, crowded: stem
forked: spike 4-slowered.

With. Arr. 212.

Lesser Water-Caltrops. Frog's Lettuce. Ditches, flow streams. June.

P. Leaves spear-shaped, alternate, cris pumo or opposite, waved and serrated.

With Arr. 213. Curt. 5.

Bloff. white, or reddish.

Greater Water-Caltrops. Curled Tet. Tet. Pondweed. Ponds, flow fireams.

June.

P. Leaves briftle-shaped, parallel, pectinatum. near together, pointing 2 ways.

With. Arr. 213. E. Bot. 323.

Bloff. whitish.

Fennel-leaved Pondweed. Rivers, Tet. Tet. ponds. July.

SAGI'NA. See Gen. Ch. p. 37.

Stem rather upright, pubefcent: apet ala. flowers alternate: without petals: petals very minute: notched at the end.

With. Arr. 216. Curt. 5.

Tet. Tet. Annual Pearlwort. Walls, gravel-walks, dry places.

> Bloff. falver-shaped, MYOSO'TIS. 5-cleft, lobes notched: Mouth closed with projecting scales: Nuts 4, imperforated.

palúftris. M. Seeds fmooth: calyxes inverfely egg-shaped, blunt, smooth, as long as the tube of the bloffom: leaves spear-shaped.

With. Arr. 225, Curt. 3.

L

Bloff. fine blue, valves forming a bright yellow eye.

Water Moufe-ear. Scorpion-grafs. Myofotis. Scorpioides. Wet ditches, springs. July.

M. Seeds fmooth: calyxes eggshaped, tapering to a point, very hairy, longer than the tube of

Pen. Mo.

arven fis.

the bloffom: leaves egg-fpearfhaped.

With. Arr. 225.

A fmaller plant than the preceding.

Field Moufe-ear. Scorpion-grass. Pen Mo. Myosotis Scorpioides. June. Walls, dry pastures, gardens.

LITHOSPER'MUM. See Gen. Ch. p. 37.

L. Seeds fmooth: bloffoms hardly officinále. longer than the cup: leaves spear-fhaped.

With. Arr. 226. E. Bot. 134.

Woodv. 213.

Bloff. pale yellow.

1,

th:

e-

35

m:

. 3.

ga

ass.

Vet

uly.

gg-

very

e of

Gromill. Graymill. June. Pen. Mo. Woods, in a dry gravelly foil.

L. Seeds wrinkled: bloffoms hard- arven'fe. ly longer than the cups.

With. Arr. 226. E. Bot. 123.

Bloff. white. June.

Corn Gromwell. Baftard Alkanet. Pen. Mo. Corn-fields.

ANCHU'SA. Bloff. funnel-shaped:
Mouth closed by projecting valves:
Nuts 4, each of one cell: perforated at the base.

femper-virens. A. Fruit-stalks in heads, 2 leaves to each head: leaves egg-shaped, acute: floral leaves somewhat waved and serrated.

With. Arr. 227. E. Bot. 45.

Bloff. fine blue.

Pen. Mo.

Evergreen Alkanet. July. Road fides, rubbish.

CYNOGLOSSUM. Bloffom funnelfhaped: Mouth closed by projecting valves: Nuts 4, depressed, fixed to the style by the inner side only, imperforated.

fylvat'icum.

C. Stamens shorter than the bloffom: leaves spear-shaped, harsh, flat, distant, nearly sitting.

With. Arr. 228.

S.

Flowers dull blue, or reddish.

Pen. Mo. Hounds-tongue. Shady lanes.

PULMONA'RIA. See Gen. Ch. p. 19.

P. Root-leaves spear-shaped. angustisólia. With. Arr. 228.

Bloff. red at first, expanding, soon changing to blue.

Bugloss Cowslips. Narrow-leaved Pen. Mo. Lungwort. Woods.

SYM PHYTUM. Bloff. funnel-shaped, bellying towards the top: Mouth closed by hollow radiate valves, which have an open hole on the outside near the borders: Nuts 4, perforated.

Leaves egg-shaped, decurrent. officinále.
 With. Arr. 230. Curt. 4. Woodv.
 215.

Bloff. yellow white.

t

Ţ.

1-

0-

d,

de

of-

ſh,

28.

19.

Common Comfrey. Banks of ri- Pen. Mo. vers, wet ditches.

Var. 2. Red-flowered.

 Leaves half decurrent; the up-tuberófum. permoft oppofite.

With. Arr. 230.

Bloff. yellow white.

Tuberous - rooted Comfrey. Fen Pen. Mo. banks.

PRI'MULA. See Gen. Ch. p. 1.

farinófa. P. Leaves scolloped, smooth: border of the blossom flat.

With. Arr. 235. E. Bot. 6.

Bloff. bluish red.

Pen. Mo. Bird's-eye Primrofe. Marshes, bogs, mountains.

LYSIMA CHIA. Bloff. wheel-shaped: Caps. globular, dagger-pointed: of 1 cell, 5 valves, and many seeds.

nem orum. L. Leaves egg - shaped, acute:
flowers solitary: stem trailing.

With. Arr. 237. Curt. 5.

Bloff. yellow, fmaller than the leaves.

Pen. Mo. Yellow Pimpernel of the Woods.

Moist shady places. June.

Anaga'llis. Bloff. wheel-shaped: Caps. cut round; of 1 cell, and many seeds.

arven's. A. Leaves egg-spear-shaped: stem trailing: calyx segments spear-shaped.

With. Arr. 238. Curt. 1.

Bloff. scarlet.

Male Pimpernel. Common Pimpernel. Corn - fields, fandy Pen. Mo. places. Aug.

Var. 2. Bloff. blue.

CAMPA'NULA. Bloff. bell-shaped: Filaments broad and arched at the base: Summit 3-cleft: Caps. beneath; 3-celled, opening at 3 lateral holes.

C. Leaves heart-shaped, 5-lobed, hederácea. on leaf-stalks, smooth; stem feeble.

With. Arr. 244. E. Bot. 73.

Bloff. pale blue.

d

:

5. he

ds.

ne.

d:

ind

em

ear-

1. 1.

Ivy-leaved Bell-flower. Moist Pen Mo. shady places. Aug.

LONICE'RA. See Gen. Ch. page 6.

L. Fruit-stalks 2-flowered: berries Xylosteum. distinct; leaves very entire, pubescent.

With. Arr. 247.

Bloff. yellow.

Upright Honeyfuckle.

Pen. Mo.

Euo'nymus. Bloff. 5 petals: Capf. coloured; 5-fided, 5-celled, 5-valved: feed-coat hollow: Seed veiled.

europæ'us. E. Flowers mostly 4-cleft: leaves fitting.

With. Arr. 259. E. Bot. 362.

Petals greenish white.

Pen Mo. Common Spindle Tree. Prickwood.
Woods, hedges. June.

Vi'ola. See Gen. Ch. page 6.

V. Stem unbranched, upright, 3cornered: leaves fringed with hairs, lower ones heart-shaped, upper ones spear-shaped.

With. Arr. 263.

Bloff. pale, bright yellow.

Pen. Mo. Yellow Violet, or Pansies. Mountainous pastures. Sept.

RI'BES. See Gen. Ch. page 40.

rúbrum. R. Bunches fmooth, pendant: flowers flattish.

With. Arr. 264. Woodv. 74.

Bloff. greenish white. Berries red.

Red Currants. Woods in the northern counties.

Var. 2. Sweet Currants.

Var. 3. Small Currents.

Pen. Mo.

Wimbledon Park, Surry.

R. Bunches upright: floral-leaves alpinum. longer than the flowers.

With. Arr. 264.

Sweet Mountain Currants. Woods Pen. Mo. and hedges in Yorkshire.

R. Bunches hairy: flowers oblong. nigrum. With. Arr. 265. Woodv. 75.

Squinancy Berries. Black Currants. Pen. Mo. Wet hedges, banks of rivers.

VIN'CA. See Gen. Ch. page 18.

V. Stems upright: leaves egg- májor. shaped: flowers on fruit-stalks.

With. Arr. 269. Curt. 4.

Bloff. blue, with a blush of purple.

Greater Perriromkle. Woods, Pen. Mo. hedges.

CHENOPO'DIUM. Cal. with 5 clefts

and 5 ribs: Bloff. none: Seed 1, round, but flatted, superior, horizontal, covered by the closing calyx.

Bonus-Henricus. C. Leaves triangular-arrow-shaped; very entire: spikes compound, leasless: axillary.

With. Arr. 270. Curt. 3.

Pen. Di.

Bloff. greenish white.

Good King Henry. Road fides.

HYDROCO TYLE. Umbel fimple, on a foot-stalk: Involucre of 2 or 4 leaves: Petals entire: Fruit compressed, bulging, divisible into two parts.

vulgáris.

H. Leaves target-shaped, notched: umbels 5-flowered.

With. Arr. 284. Curt. 6.

Flowers reddish white.

Pen. Di.

Marsh Pennywort. White Rot. Marshy ground. June.

SANICULA. Umbellules crowded, forming a kind of head: Florets of the centre barren: Fruit set with hooked prickles.

S. Root-leaves fimple: florets fit- europæ'a. ting.

With. Arr. 284. E. Bot. 98.

Bloff. white.

Common Sanicle. Woods, thickets. Pen. Di. June.

CAU'CALIS. Bloff. radiated: florets of the centre male: Involucre leafits entire: Petals bent inwards, notched at the end: Seeds covered with prickles or briftles.

C. Involucellums extending half daucoides, way round, strap shaped: seeds egg-oblong: angles on the back 4, distant, rough with bristles: leaves smooth, hairy on the ribs and at the base.

With. Arr. 287. E. Bot. 197.

Florets pink.

Fine-leaved Baftard Parfley. Cal- Pen. Di. careous corn-fields. June.

C. Seeds egg-shaped, tapering to scandicina a point; with short hooked prickles on the back.

With. Arr. 289. Curt. 1.

Petals white.

Pen. Di. Scandix Anthrifcus. Rubbish, road sides. June.

Bu'nium. Bloff: uniform: Umbel crowded: Styles bent back deciduous: Seeds rather cylindrical, fcored, thicker towards the end.

deciduous: stem leasses at the base, tapering downwards, zigzag: styles permanent.

Pen. Di. With Arr 201 Curt 4.

With. Arr. 291. Curt. 4. Earth Nut. Pig Nut. June.

Æтни's A. Involucellums reaching half way round, 3 leaves bent downwards: Fruit nearly globular, deeply furrowed.

Æ. All the leaves divided into many briftle-fhaped fegments: involucrum 1 leaf: fruit eggoblong, tapering at the end.

With. Arr. 305.

Bald, or Bawd Money. Mountainous pastures.

Méum.

Pen. Di.

Scan'dix. Bloff. radiated, central: Florets frequently male: Petals notched at the end: Styles permanent: Fruit awl-shaped.

S. Seeds gloffy, cylindrical, and Cerefelium. beaked: umbels lateral, nearly fitting.

With. Arr. 307.

Bloff. white.

Common Chervil. Shepherd's Needle. Pen. Di. Hedges.

CHEROPHY LLUM. Involucellum reflected, concave: Petals heartfhaped, bent inwards: Fruit shining, generally smooth, oblong.

C. Stem smoothish, scored, a little sylvestre. swoln at the knots.

With. Arr. 308. Curt. 4.

Stem. woolly. Flowers white.
Wild Cicely. Cow Parfley. June. Pen. Di.
Hedges, orchards, pastures.

SMYR'NIUM. Petals keeled, tapering to a point: Fruit egg-globular, bulging, angular with ribs. Olufátrum.

S. Stem-leaves growing by threes; on leaf-stalks, ferrated.

With. Arr. 310. E. Bot. 330.

Pen. Di.

Flowers greenish yellow.

Alifanders. Alexanders. Ditches, rocks on the fea coaft.

CA'RUM. Involucre 1 leaf: Petals united, bent inwards, notched at the end: Fruit small, elliptical, bulging, roundish and scored.

Cárui.

C.

With. Arr. 311. Woodv. 45.

Pen. Di.

Petals tinged with red. June. Common Carraway. Meadows, pastures.

PIMPINEL'LA. Petals bent inwards: Styles upright: Summits nearly globular: Fruit small, egg-oblong: with 5 elevated ridges.

dioica.

P. Plant dwarfish: umbels very numerous, compound and simple: male and hermaphrodite flowers on distinct plants.

With. Arr. 313.

Flowers yellowish, or whitish.

Least Anise. Mountains, pastures. Pen. Di.
June.

ÆGOPO DIUM. Fruit egg-oblong, fcored, tapering at each end.

Æ. Upper leaves 3 together, lower Podagrária. ones in triple threes.

With. Arr. 314.

Bloff. white.

Herb Gerard. Ground Ash.

Pen. Di.

Orchards, gardens, pastures.

GENTIA'NA. Bloff. 1 petal: Recept. of the feeds 2, placed lengthways: Capf. 2-valved, 1-celled.

G. Bloffom bell-shaped, not beard-collina. ed, pistil woolly: summits 3: leaves strap-spear-shaped, hairy.

With, Arr. 282.

Gentian. Aften Down.

Pen. Di.

VIBUR'NUM. Cal. with 5 divisions; fuperior: Bloff. 5-cleft: Berry of 1 cell; closed: Seed 1.

96	MAY.
Lantána.	V. Leaves heart-shaped, serrated, veined, cottony underneath. With. Arr. 315. E. Bot. 331.
	Bloff. white. Berries black.
Pen. Tri.	Pliant Mealy Tree. Wayfaring Tree. Woods in a calcareous foil.
ópulus.	V. Leaves lobed: leaf-stalks fet with glands. With. Arr. 315. E. Bot. 332.
	Bloff. white. Berries red.

Water Elder. Guelder-rofe. Pen. Tri. Woods, wet hedges. June.

> STATICE. Cal. 1 leaf, entire, plaited, dry, permanent: Petals 5: Capf. 1-celled, without valves: Seed upright.

S. Stalk fimple, with a head of Arméria. flowers: leaves strap-shaped. With. Arr. 320. E. Bot. 226. Bloff. pink. June. Pen. Pen. Common Thrift. Sea Gilliflowers.

> LI'NUM. Cal. 5-leaved: Petals 5: Capf. 5-valved, 10-celled: Seeds folitary.

L. Leaves opposite, egg-spear- cathar ticum shaped: stem forked: blossom pointed.

With. Arr. 323. Curt. 3.

Purging Flax. Mill Mountain. Pen. Pen. Pen. Dry meadows, pastures. July.

Myosu'Rus. Cal. 5 leaves, united at the base, deciduous: Nectaries 5, like petals; awl-shaped: Seeds numerous, coated, pendent.

M. With. Arr. 326. Curt. 4. min'imus.

Flowers yellowish green.

Little Mouse-tail. June. Pen. Poly. Corn-fields, meadows, in gravel-

forn-fields, meadows, in gravelly foil.

LEUCO'JUM. Bloff. bell - shaped, with 6 divisions, thickest at the ends: Summit undivided.

L. Sheath many-flowered: ftyle Æftívum. club-shaped.

With. Arr. 331. Curt. 5.

Flowers white, streaked. Hex. Mo. Thames side.

NARCIS'SUS. See Gen. Ch. p. 12.

poéticus:

N. Sheath 1-flowered; nectary wheel-shaped, very short, skinlike, finely scolloped.

With. Arr. 331. E. Bot. 275.

Hex. Mo.

Flower folitary, pure white. Sandy heathy places.

biflorus.

N. Sheath 2-flowered: nectary wheel-shaped, very short, skin-like, finely scolloped: leaves acute on the keel, the edges turned inwards.

With. Arr. 332. E. Bot. 276.

Bloff. pale whitish cream-colour.

Hex. Mo.

Pale Daffodil. Meadows, hedges. Rare.

AL'LIUM. Bloff. 6 petals, expanding: Sheath dry and skin-like, many-flowered: umbel crowded: Caps. superior, 3-celled.

ursinum.

A. Stalk 3 fquare: leaves spearshaped, on leaf-stalks: umbel flattopped.

With. Arr. 334. E. Bot. 122.

Flowers large, numerous, white.

Ramfon Garlic. Ramfons. Woods, Hex. Mo. hedges, dry meadows.

SCIL'LA. See Gen. Ch. p. 20.

S. Bunch of few flowers, furnished ver na. with floral leaves: bloffoms bell-shaped: leaves from the root, strap-shaped channelled: bulb coated.

With. Art. 338. E. Bot. 23.

Bloff. blue.

Vernal Squill. June. Hex. Mo. Meadows, pastures.

CONVALLA RIA. Bloff. 6-cleft: Berry 3-celled, superior: Seeds 2.

C. Stalk naked, femi-cylindrical: majális. flowers spiked, nodding.

With. Arr. 341. Curt. 5.

Bloff. white, highly fragrant.

Lily of the Valley. May Liby. Hex. Mo.

Woods, heaths, near Hampstead.

C. Leaves alternate, embracing Polygonathe stem: stem 2-edged: fruittum.

MAY.

ftalk axillary, mostly 1-flowered.

With. Arr. 342. E. Bot. 280. Woodv. 44.

Hex. Mo. Sweet-smelling Solomon's Seal.

Fiffures of rocks. June.

multiflóra. C. Leaves alternate, embracing the stem: stem cylindrical: fruitstalks axillary, many-flowered.

With. Arr. 342. E. Bot. 279.

Woodv. 44.

Flowers white tipped with green. Common Solomon's Seal. Woods.

June.

Hex. Mo. Var. 2. Dwarf English Solomon's Seal. Woods in Wiltshire.

A conus. Spadix cylindrical, covered with florets: Bloff. 6 petals, naked: Style none: Capf. 3 celled.

cálamus. A. Floral-leaf very much longer than the spike.

With. Arr. 343. E. Bot. 356. Woodv. 173.

Hex. Mo. Myrtle Flag. Sweet-smelling Flag. Banks of rivers. June. BER'BERIS. Cal. 6-leaved: Petals 6, with 2 glands to the claw: Style none: Berry superior, 1-celled, open at the end: Seeds 2 or 3.

B. Fruit-stalks forming bunches: vulgáris. thorns 3 together. June.

With. Arr. 350. E. Bot. 49.

Woodv. 234.

Bloff. yellow. Berries red. Hex. Mo. Barberry. Pipperidge-bush.

RU'MEX. Cal. 3-leaved: Petals 3, closing: Seed 1, three-cornered, enclosed in the blossom.

R. Styles 2. With. Arr. 357.

Mountain Dock. Welsh Sorrel. Hex. Tri. Mountains. July.

R. Leaves fpear-halberd-shaped. Acetosel'la. With. Arr. 358. Curt. 5.

Flowers reddifh.

Sheep's Sorrel, or Dock. Sandy Hex. Tri. meadows. June.

TRIGLO CHIN. Cal. 3-leaved: Petals 3, like a cup: Style none: Capf. opening at the base. Aug. maritimum T. Capfule 6-celled, egg-shaped.

With. Arr. 359. E. Bot. 255.

Sea Arrow-grass. Meadows near
the sea.

Poly gonum. See Gen. Ch. p. 45.

Bistor'ta. P. Leaves egg-shaped, extending at the base along the leaf-stalks.

With. Arr. 382. Curt. 1. Woodv.

34.

Oct. Tri. Bloff. with 5 divisions, pale red.

Great Bistort. Snake-weed. June.

PA'RIS. Cal. 4-leaved: Petals 4, narrower: Berry 4-celled.

quadrifólia. P. Stem leaves generally 4.

With. Arr. 385. E. Bot. 7.

Bloff. pale green. June.

Oct. Tet. Herb Paris. True-love. Woods, thady places.

MYRIOPHYL'LUM. Male and Fem. flowers frequently on the same plant. Cal. 1 leaf, 4-toothed: Petals 4, with claws: soon falling off: Styles none: Nuts 2 to 4, covered with a coat.

M. Spikes interrupted, leaflefs. spicatum. With. Arr. 389 E. Bot. 83.

Flowers 4 in a whirl.

Spiked Water Milfoil. Ditches, Oct. Tet. ponds. July.

AR BUTUS. Cal. with 5 divisions: Bloff. egg-shaped, pellucid at the base: Mouth 5-clest: Berry 5-celled, many-seeded; superior.

A. Stems trailing: leaves wrinkled, alpina. formewhat ferrated, and fringed with hairs.

With. Arr. 399.

Berries black, globular.

Mountain Strawberry Tree. Dry Dec. Mo. mountains in Scotland.

A. Stems trailing: leaves oblong Uva-ur'fi. egg-shaped, very entire, veined like network underneath.

With Arr. 399. Woodv. 70. Dec. Mo. Flowers flesh colour. Berries red.

June.

SAXTERAGA. See Gen. Ch. p. 46.

S. Stem-leaves strap-shaped, en- hypnoides.

Dec. Tri.

tire, or 3-cleft; fuckers trailing: flem with few leaves.

With. Arr. 407.

Dec. Di. Petals greenish white. July.

Ladies Cushion. Moss Saxifrage.

SILE'NE: See Gen. Ch. p. 48.

notched at the end: capfules globular.

With, Arr. 416.

Flowers fingle, bright purple.

Moss Catchfly. Mountains. June.

ARENA'RIA. Cal. 5 leaves, expanding: Petals 5, entire: Capf. 1-celled, many-feeded.

triner'via. A. Leaves egg-shaped, 3-fibred, pointed, on leaf-stalks.

With. Arr. 421. Curt. 4.

Petals white.

Plantain-leaved Sandwort.

Woods, wet hedges. July.

ferpyllifilia. A. Leaves fomewhat egg-shaped, acute, sitting: blossoms shorter than the calyx.

With, Arr. 421. Curt. 4.

Flowers white.

July. Dec. Tri.

Least Chickweed. Thyme-leaved Sandwort.

A. Leaves femi-cylindrical, taper-marina, ing, opposite, as long as the joints: stipulæ membranaceous, acute, stems prostrate: capsules longer than the calyx.

With. Arr. 422.

Flowers purplish.

Dec. Tri.

Salt marshes.

Oa.

A. Leaves awl-shaped: stems pa-ver'na. nicled: calyx taper - pointed, scored.

With Arr. 423.

Mountain Sandwort. Mountain- Dec. Trious fituations. Aug.

STELLA'RIA. See Gen. Ch. p. 23.

S. Leaves strap-shaped, very en- gramin eastire: flowers in panicles.

With. Arr. 419.

Petals white.

Dec. Tri.

Lesser Stichwort. Meadows, pastures, hedge-banks. SE'DUM. Cal. 5-cleft: Bloff. none, or 5 petals: Neclariferous Scales 5, at the base of the germens: Caps. 5 distinct, like a legumen.

ang licum. S. Leaves somewhat egg-shaped, growing to and sitting, bulging, alternate: tust branched, forked.

With. Arr. 428. E Bot. 171.

Bloss white, or purplish white.

English States grow. Books roofs.

Dec. Pen. English Stone-crop. Rocks, roofs, walls. June.

Ox'ALIS. Cal. 5 leaves: Petals connected by the claws: Capf. 5-fided, opening at the corners.

corniculáta. O. Stem reclining, herbaceous : fruit-ftalks forming umbels. With. Arr. 430.

Bloff. yellow.

Dec. Pen. Yellow flowered Cuckow-bread.

Near Exeter. Oct.

LYCH'NIS. Cal. 1 leaf, oblong, even: Petals 5, with claws; limbs often cloven: Capf. 1 to 5-celled.

L. Stamens and piftils in diftinct dioica.
plants: capfules 1-celled.

Var. 1. diur na. Bloff. red. With. Arr. 431. Curt. 2.

Red-flowered Wild Campion, or Cuckow-flower. Sept. Dec. Pen. Hedges, ditches, moist woods.

L. Petals nearly entire. visca'ria. With, Arr. 433.

Red German Catchfly. June. Dec. Pen. Mountainous pastures.

CERASTIUM. See Gen. Ch. p. 12.

C. Leaves egg-fhaped: petals as vulgátum. long as the calyx: ftems fpreading.

With. Arr. 433. Curt. 2.

Petals white. Aug.

Narrow-leaved Mouse-ear. Mea- Dec. Pen. dows, pastures, walls.

C. Leaves strap-spear-shaped, blunt- arvense. ish, fringed at the base: blossoms larger than the calyx.

With.Arr. 434. Curt. 6. E. Bot. 93.

Corn Mouse-ear. Sept. Dec. Pen.

Corn-fields, gravelly meadows.

As ARUM. Cal. 3 or 4-cleft, fitting on the germens: Bloff. none: Capf. like leather, crowned, of 6 cells: but no valves.

europæ'um. A. Leaves kidney-shaped, blunt, in pairs.

With. Arr. 440. Woodv. 86.

Flowers purplish.

Dode. Mo. Afarabacca. Woods, shady places,

CARPI'NUS. Male and Fem. flowers on the same plant: Bloff. none: Cal. 1 leaf, a fringed scale.

Male. Stamens 18 or 20.

Fem. Germens 2, with 2 styles on each: Nut egg-shaped.

Bet'ulus. C. Scales of the cones flat.

With. Arr. 442.

Bark fmooth, white.

Dode. Di. Common Hornbeam Tree. Woods, hedges.

Fa'gus. See Gen. Ch. p. 24.

F. Leaves spear-shaped, with tapering serratures, naked underneath.

With. Arr. 443.

Chefnut Beech. Chefnut Tree. Dode. Di. Woods, hedges.

EUPHOR BIA. Cal. 1 leaf, distended: Bloff. 4 or 5 petals, fitting on the calyx: Capf. 3 united.

E. Umbel with generally 5 spokes: Parális. spokes cloven: involucellums heart-kidney-shaped: leaves tiled upwards.

With. Arr. 448. E. Bot. 195.

Plant sea green. Sept. Sea Spurge. Sea shores.

E. Umbel with many fpokes; amygdafpokes forked: involucellums perforated, circular: leaves blunt.

With. Arr. 452. E Bot. 256.

Wood Spurge. Woods, hedges.

E. Umbel with 5 spokes; spokes portlan'dica. forked: involucellums somewhat heart-shaped concave: leaves

ftrap-fpear-shaped, smooth, expanding.

With. Arr. 448.

Flowers nearly fitting, yellow.

Dode. Tri. Portland Spurge. Sandy fea-shores.
Aug.

Cyparis's E: Umbel with many spokes; spokes forked: involucellums nearly heart-shaped: leaves on the stem spear-shaped, on the barren branches bristle-shaped.

With. Arr. 451.

Dode. Tri. Woods at Enville. June.

PRU'NUS. See Gen. Ch. p. 25.

Pádus. P. Flowers in bunches: leaves deciduous, with 2 glands at the base on the under side.

With. Arr. 455.

Ico. Mo. Birds' Cherry. Wild Cluster Cherry. Woods, hedges.

fpear-shaped, downy underneath, doubled together.

With. Arr. 456.

dco. Mo. Black Cherry Tree. Mazzards.

Hedges in Suffolk.

CRATE GUS. See Gen. Ch. p. 50.

C. Leaves egg-shaped, cut, fer- A'ria. rated, cottony underneath.

With. Arr. 458.

Petals white.

.

3.

S.

S

e

.

White-beam Hawthorn. Woods, Ico. Di. hedges.

C. Leaves heart shaped, with 7 torminális. angles: lowermost lobes strad-dling.

With. Arr. 458. E. Bot. 298.

Bloff. white.

Wild Service Tree, or Sorb. Ico. Di.

Woods, hedges.

Son'sus. See Gen. Ch. p. 50.

 Leaves wing-cleft, cottony un- hy brida. derneath.

With. Ann. 461.

Baftard Service. Mountains. Ico. Tri.

Mes'PILUS. Cal. 5-cleft: Petals 5: Berry with 1, 2, or 5 cells.

M. Thornless: leaves spear-shap- germanica.

ed, cottony underneath: flowers folitary, fitting.

With. Arr. 461.

Bloff. white.

Ico. Pen. Medlar Tree. Hedges.

> Py Rus. See Gen. Ch. p. 51.

P. Leaves ferrated: flowers in um-Málus. bels, fitting.

With. Arr. 462. E. Bot. 179.

Ico. Pen. Crab Tree. Wilding. Woods, hedges.

> Ru'Bus. Cal. 5-cleft; Petals 5: Styles from the top of the germens: Drupa cluftered, 1-celled, fixed to a conical receptacle fo as to refemble a berry.

1dæ us. R. Leaves winged, with 5 or 3 leafits: stem prickly: leaf-stalk channelled.

With, Arr. 468. Woodv. 138.

Bloff. white. Berry red. Rasp-berry Bush. Bramble. June.

Woods, hedges.

Ico. Poly.

R. Leaves 3 together: ftem with- ar'cticus. out prickles, supporting a fingle flower.

With. Arr. 470.

Flowers purplish. Berry dark Ico. Poly. purple.

Highlands of Scotland.

R. Leaves fimple lobed: ftem Chamæwithout prickles, with 1 flower: morus. male and female flowers on different plants.

With. Arr. 471.

Bloff. white, or purple. Ico. Poly. Cloud-berry. Bramble. June.

- TORMENTIL'LA. Cal. 8-cleft: Petals 4: Seeds roundish, naked, fixed to a small juiceless receptacle.
- T. Stem trailing: leaves on leaf- rep'tans. ftalks.

With. Arr. 476.

Crecping Tormentil. July. Ico. Poly. Woods, fandy barren pastures.

FRAGA'RIA. See Gen. Ch. p. 13.

MAY.

vésca.

F. Leaves 3 together: runners creeping.

With. Arr. 471.

Bloff. white. Fruit red.

Ico. Poly. Woods, hollow ways.

Tune.

DRY'AS. Cal. 5 or 10-cleft: Petals 5 or 8: Seeds with tails, formed by the feathered ftyle: Receptacle broad and flat.

octopetala. D. Petals 8: leaves fimple.

With. Arr. 478.

Bloff. fnowy white.

Ico. Poly.

Mountain Avens. High mountains. June.

GEUM. Cal. 10-cleft: Petals 5: Styles terminating: Seed with a jointed awn: Recept. pillar-like.

urbánum.

G. Flowers upright: fruit globular, woolly: awns hooked, bare: root-leaves lyre-shaped: stemleaves in threes.

With. Arr. 477. Curt. 2. Woodv. 259.

Petals yellow.

Common Avens. Herb Bennet. Ico. Poly. Woods, hedges. Aug.

ACTE'A. Bloff. 4 petals: Cal. 4 leaves: Berry 1-celled: Seeds femi-circular.

A. Bunch egg-shaped: fruit berry- spicáta. like.

With. Arr. 483.

Bloff. white. Berries black.

Herb Christopher. Baneberries. Poly. Mo.
Woods, shady places. June.

CHELIDO'NIUM. Bloff. 4 petals: Cal. 2 leaves: Pod strap-shaped: Receptacle of the seeds (generally) lattice-like.

C. Fruit-stalks forming umbels. május. With. Arr. 483. Woodv. 263.

Flowers yellow.

a

Greater, or Common Celandine. Poly. Mo. Hedges, shady places. July.

C. Fruit-stalks with 1 flower: hy'bridum. leaves wing-cleft: segments strap-

fhaped: ftem fmooth: pod with 3 valves.

With. Arr. 485. E. Bot. 201.

Poly. Mo. Plant sea green. June.

Red Celandine. Sandy cornfields.

Cis'tus. Bloff. 5 petals: Cal. 5 leaves; 2 of them smaller: Caps. from 1 to 10-celled: from 3 to 10-valved.

ang'licus. C. Stems trailing; leaves hairy, opposite, oblong, edges rolled back: flowers in bunches.

With. Arr. 490.

Petals yellow, finely fcolloped.

Poy. Mo. Hairy Ciftus. Mountainous paftures, rocks.

Helian themum. leaves oblong, edges rolled back, fomewhat hairy.

With. Arr. 492. Curt. 5.

Petals yellow, toothed on the outer part.

Dwarf Ciftus. Mountainous meadows, in a calcareous foil.

Aug.

Var. 2. Bloff. white.

Poly. Mo.

Var. 3. Bloff. rose-coloured.

AQUILE GIA. Cal. none: Petals 5: Nectaries 5: horn-shaped, alternating with the petals: Caps. 5, distinct.

A. Nectaries bowed inwards, near-vulgáris. ly equal to the petals: leafits on leaf-stalks, lobes distant, roundish, bluntish.

With. Arr. 495. E. Bot. 297.

Bloff. blue or purple: pendent.

Columbines. Woods, thickets. June.

Poly. Pen.

A'RUM. Sheath 1 leaf, cone-shaped: Fruit naked above, bearing germens at its bottom, and stamens in the middle.

A. Leaves halberd - shaped, very miculátume entire: spike-stalk club-shaped.

With. Arr. 497. Curt. 2. Woodv.

25.

Spike-flalk, upper part purple, or buff. Berries red.

Poly. Poly. Cuckoo-pint. Wake Robin. Lords and Ladies. Shady places, ditch banks.

ANEMO'NE. See Gen. Ch. p. 25.

ranunculoídes. A. Seeds pointed: leafits fnipt: petals roundish: stem mostly 2-flowered.

With. Arr. 499.

Var. 2. Stem zig-zag. Bloff. 5 petals.

Poly. Poly. Yellow Anemone. Shady places, hedges.

ADO'NIS. Cal. 5 leaves: Petals 5 or more: Nect. none: Seeds naked.

æstivális. A. Flowers with 5 petals: fruit egg-shaped.

Poly. Poly. Corn-fields on Salifbury Plain.

RANUN'CULUS. See Gen. Ch. p. 7.

fcelerátus. R. Lower leaves hand-fhaped, the upper fingered: fruit oblong.

With. Arr. 505. Curt. 2.

Flowers small yellow.

Round - leaved Water Crowfoot.

Celery-leaved Crowfoot. Shal- Poly. Poly. low waters.

June.

R. Seeds rough, with tubercles parviflórus. ending in hooked points; leaves heart-shaped, hairy, lobed, or toothed: stem prostrate.

With, Arr. 506. E. Bot. 120.

Flowers small yellow.

Small flowered Crowfoot. Corn- Poly. Poly. fields, in a gravelly foil.

R. Leaves under water hair-like; aquatilisthose above with nearly central leaf-stalks.

With. Arr. 507. E. Bot. 101.

Flowers white, with a yellow spot at the base. July. Poly. Poly. Water Crowfoot. Ponds, ditches.

R. Root bulbous: calyx reflected: bulbósus. fruit-stalks furrowed: stem upright, many-flowered: leaves compound.

With. Arr. 508. Curt. L.

Poly. Poly. Petals bright yellow.

Butter-flower: Butter-cups. Meadows.

TROL'LIUS. Cal. none: Petals about 14: Capf. many, egg-shaped, many-feeded.

europæ'us. T. Petals approaching: nectaries as long as the stamens.

With. Arr. 510. E. Bot. 28.

Bloff. closed, yellow.

Poly. Poly. Globe-flower. June. Sides of mountains.

A'juga. See Gen. Ch. p. 54.

rep'tans. A. Plant fmooth, with creeping fuckers: leaves egg-fhaped, fcolloped.

With. Arr. 517. Curt. 2.

Bloff. blue, red, or white.

Didy. Gym. Common Bugle. July.

Moift meadows, pastures, woods.

GALEOB'DOLON. Bloff. upper-lip entire, vaulted: lower lip without teeth, 3-cleft; fegments broad, pointed: Anthers fleshy on the back.

G. lúteum.

With. Arr. 530. Curt. 4.

Bloff. yellow.

Yellow Archangel, or Dead Nettle. Didy. Gym. Woods, shady places.

SCROPHULA RIA. Cal. 5-cleft: Bloff. with 5 divisions: tube globular, the lower segment reflected: Caps. 2-celled, partition double.

S. Leaves oblong heart-shaped, 3- nodósa. fibred at the base, corners of the stem acute.

With. Arr. 553.

Bloff. upper segments purple.

Great Figwort, Kernelwort. Didy. Ang. Woods, moist hedges. July.

DIGITA'LIS. Cal. with 5 divisions: Bloff. beil-shaped: 5-cleft, bellying: Capf. egg-shaped, 2-celled, many-seeded.

D. Segments of the calyx egg- purpurea.

fhaped, acute: bloffom blunt, upper lip nearly entire.

With. Arr. 555. Curt. 1. Woodv. 24.

Bloff: purple elegantly mottled within.

Common Foxglove. July.

Didy. Ang. Hedge-banks; fides of hills in gravelly or fandy foil.

OROBAN CHE. Cal. fomewhat cloven: Bloff. gaping: Germen refting on a gland: Capf. 1-celled, 2-valved, many-feeded.

májor. O. Stem unbranched, pubescent: stamens the length of the blosfom.

With. Arr. 557. Curt. 4.

Flowers ruft colour, purplish.

Didy. Ang. Common Broomrape. June.

On roots of trees, and in cornfields.

MOEN'CHIA. Pouch entire, eggfhaped, crowned with the ftyle: Valves rather convex, parallel to the partition: Cells with many feeds.

M. Pouches inverfely egg-shaped, fativ'a. on fruit-stalks, containing many feeds.

With. Arr. 562.

Bloff. yellow. June. Tetd. Silicu.

Gold of Pleasure. Common Camline.

CRAM'BE. The 4 longer filaments cloven at the top, one of the clefts bearing the anther: Seed-vessel simple, globular, deciduous.

C. Leaves and flem fmooth. maritima. With, Arr. 563.

Bloff. large, white.

Sea Colewort. Cliff Kale. Sandy Tetd. Silicustea fea shores. June.

DRA'BA. See Gen. Ch. p. 27.

D. Leaves hairy, toothed, or entire: ftaff with 1 leaf: pouches
oblong, upright: petals entire.

With. Arr. 565.

MAY.

Tetd. Silieu. Petals pure white.

Highlands of Scotland. June.

murális. D. Stem branched: leaves eggshaped, fitting, toothed.

With. Arr. 566.

Tetd. Silicu. Flowers white.

Speedwell-leaved Whitlow-grafs.

D. Stem-leaves numerous, hoary:
pouches oblong, flanting, nearly
fitting.

With. Arr. 566. E. Bot. 388.

Petals white, flightly notched.

Tetd. Silicu. Wreathen-podded Whitlow-grafs.

Fiffures of limestone rocks.

July.

THLAS'PI. See Gen. Ch. p. 26.

campéstre. T. Pouches roundish: leaves arrow-shaped, toothed, hoary.

With. Arr. 569. Curt. 5.

Bloff. white. July.

Tetd. Silicu. Mithridate Mustard. Bastard Cress. Corn-fields, in clayey fandy soil.

- COCHLEA'RIA. Pouch notched at the end, turgid, rough, manyfeeded: valves bulging, blunt.
- C. All the leaves trowel-shaped, dánica. with halberd-shaped angles at the base.

With. Arr. 573.

Bloff. white. June. Tetd. Silicu. Scurvy-grafs. Sea shores.

C. Leaves kidney-shaped, entire, groenlan-dica.

With. Arr. 537.

Mountains of Caernarvonshire. Tetd. Silicu.

C. All the leaves egg-fpear-fhap- an glica. ed.

With. Arr. 574.

Spoonwort. English Scurvy-grass. Tetd. Silicu.

C. Root-leaves spear-shaped, scol- Armorácia. loped: stem-leaves stript.

With. Arr. 575. Woodv. 150.

Var. 2. Root-leaves deeply wingcleft.

Flowers white.

Horse Radish. Sides of ditches. Tetd. Silicu.

I'BERIS. Bloff. unequal, the two outer petals larger: Pouch roundish, compressed: Partition placed crossways: 1 seed in each cell.

nudicaúlis. I. Herbaceous: leaves wing-cleft: flem naked, fimple.

With. Arr. 575. Curt. 6. E. Bot. 327.

Flowers fmall, white. July. Rock-cress. Naked Candytust.

Ted. Silicu. Rock-crefs. Naked Candytuft.

Gravelly places, barren heaths.

CARDAM INE. See Gen. Ch. p. 28.

bellidifólia. C. Leaves egg-shaped, very entire, on leaf-stalks.

With. Arr. 577.

Flowers white.

Tetd. Siliq. Daily-leaved Ladies Smock. Near Denbigh.

petrz'a. C. Leaves oblong, toothed.

With. Arr. 577.

Bloff. white. July.

Tetd. Siliq. Mountain Ladies Smock. Rocks.

C. Leaves winged: leafits fpear- impátiens, fhaped, toothed or cut: ftipulæ fringed.

With. Arr. 577. E. Bot. 80.

Petals small, white. June.

Impatient Ladies Smock. Moun- Tetd. Siliq. tainous meadows.

C. Stem zigzag: leafits toothed, flexuófa. mostly alternate, unequal at the base.

With. Arr. 578. Curt. 4.

Petals white.

Zigzag Ladies Smock. July. Tetd. Siliq. Ditches, woods, shady wet places.

Sisym'brium. Pod cylindrical, opening: valves firaightish, about the length of the partition: Cal. and Bloff. expanding.

S. Leaves notched, toothed, nak- I'ne. ed: stem smooth: pods upright.

With. Arr. 583. Curt. 5.

Bloff. fmall yellow.

Broad-leaved Water-cress. Old Tetd. Silique walls, rubbish. Aug.

ERY'SIMUM. See Gen. Ch. p. 58.

officinále. E. Pods pressed to the spike-stalk: leaves notched.

With. Arr. 583. Curt. 5. Woodv. 244.

Tetd. Siliq. Bloff. yellow. June. Hedge Mustard. Wormseed.

CHEIRAN'THUS. Germen with a glandular tooth on each fide: Cal. closed, 2 of its leafits bulging at the base: Seeds flat.

Cheiri. C. Leaves spear - shaped, acute, smooth: branches angular: stem shrub-like.

With. Arr. 586.

Flowers yellow.

Tetd. Siliq. Wall-flower. Wild Cheir. Old walls, roofs. June.

HES'PERIS. Petals turned obliquely: Glands within the shorter stamens: Cal. closed: Pod stiff and straight: Summit forked at the base, approaching at the top.

H. Stem fimple, upright: leaves inódoratoothed: petals blunt.

With. Arr. 586.

Flowers large, purplish red.

Scentles Damewort. Paftures, Tetd. Siliq. hedges. June.

A'RABIS. Nectary, glands 4, within the leafits of the calyx, refembling a reflected scale: Pod long, compressed, strap-shaped, entire, and knobbed at the end.

A. Stem-leaves spear-shaped, on thaliana, leaf-stalks, very entire: root-leaves egg-spear-shaped, toothed.

With. Arr. 587. Curt. 2.

Flowers small, white.

Turkey Pod. Codded Mouse-ear. Tetd. Siliq.

Walls, roofs, sandy places.

A. Leaves toothed, rough with firic'ta. hair: those of the root spatulashaped, those of the stem halfembracing the stem, oblong: pods 2-edged, upright.

With. Arr. 587.

Velley's Marine Plates. Shiercliff's Briftol Guide, p. 83.

Flowers in bunches, white.

Tetd. Siliq. Rough Turkey Pod. St. Vincent's Rocks near Briffol.

Turrita. A. Leaves embracing the ftem:

pods bent backwards, flat ftrapfhaped: cups fomewhat wrinkled.

With. Arr. 588. E. Bot. 178.

Flowers straw-coloured.

Tetd. Siliq. Tower Turkey Pod. June. Old walls, stony places.

TURRI'TIS. Pod very long, angular, stiff, and straight: Cal. close, upright: Bloff. upright.

glábra. T. Root-leaves toothed, rough with hair: stem-leaves very entire, embracing the stem, smooth.

With. Arr. 588. Curt. 4.

Bloff. greenish white.

Tetd. Siliq. Tower Muftard. June.

Meadows, paftures, pits.

Bras'sica. Cal. upright, close:
Glands 1 between each shorter
stamen and pistil, and 1 between
each pair of longer stamens and
the calyx: Seeds globular.

B. The root a regular continuation Nápus. of the stem: spindle-shaped.

With. Arr. 590.

Bloff. yellow.

Wild Navew. Rape. Ditch Tetd. Siliq. banks.

B. The root a regular continuation olerácea. of the stem: cylindrical sleshy.

With, Arr. 501.

Sea Colewort. Sea Cabbage. Tetd. Siliq. Cliffs on the fea coaft.

B. Leaves spear-shaped, indented-murális. serrated, smoothish: stem upright, smooth. Roughish ascending. Leaves indented or wing-cleft.

With. Arr. 592. Curt. 3.

Bloff. yellow.

Wild Rocket. Old walls, rubbish. Tetd. Siliq. June.

K 2

monen's. B. Leaves wing-cleft, wings cut, ferrated: stem nearly naked, trailing.

With. Arr. 593.

Tetd. Siliq. Isle of Man Cabbage. July. Sea shores, in fandy soil.

SINA PIS. Cal. expanding: Bloff. claws upright: Glands between the shorter stamens and the pistil, and between the longer stamens and the calyx: Pod beaked opening: valves shorter than the partition.

arven'fis. S. Pods with many angles, fwoln and bunched out by the feeds: fmooth, longer than the 2-edged beak.

With. Arr. 594. Curt. 5.

Petals yellow.

Tetd. Siliq. Chadlock. Wild Mustard, or Charlock. Corn-fields.

JUNI PERUS. Male and Fem. flowers distinct.

Male. Cal. a scale of the catkin: Bloff. none.

Fem. Cal. with 3 divisions:

Petals 3: Piftils 3: Drupa
juicy, closed, 1-celled, manyfeeded: with 3 tubercles formerly the calyx.

J. Leaves 3 together, expanding, commúnis. fharp pointed; longer than the berry.

With. Arr. 599. Woodv. 95.

Berries bluish black.
Juniper Tree. Heaths.

Mond, Tri.

GERA'NIUM. See Gen. Ch. p. 59.

G. Fruit-stalks shorter than the diffec tum, leaves: leaves 5 lobed, lobes 3-cleft, or many-cleft: petals notched, seed-coats with soft hairs.

With. Arr. 603. Curt. 6.

Flowers red.

Jagged Crane's-bill.

Aug. Mond. Dec.

Road fides, borders of fields.

G. Fruit-stalks folitary, 2-flower- phæ'um. ed, opposite the leaves: calyx

fomewhat awned: ftem upright: petals waved.

With. Arr. 605. E. Bot. 322.

Mond. Dec. Dufky Crane's bill. June.
Mountainous pastures.

rotundifólium.

G. Petals entire, as long as the calyx: ftem spreading: leaves kidney-shaped, cut: feed-coat even, hairy: feeds reticulated.

With. Arr. 607. E. Bot. 157.

Flowers purplish flesh - colour, fometimes white. July.

Mond. Dec. Round-leaved Crane's-bill. Walls, roofs, ditch-banks.

moschátum. G. Flowers in umbels: leafits mostly on short leaf-stalks unequally cut.

With. Arr. 609.

Bloff. red or purple.

Mond, Dec. Musked Crane's-bill. Dry paftures. Oct.

> Pin'us. Male and Fem. flowers on the fame plant: Bloff. none. Male. Calyx scales forming a bud standing open: Anthers naked.

Fem. Calyx scales forming a cone, 2 flowers in each scale: Pistil 1: Nut of 1 cell, without valves, bordered with a membrane.

P. Leaves in pairs, rigid: cones sylvéstris, egg-conical, mostly in pairs, as long as the leaves: scales oblong, blunt.

With. Arr. 615. Woodv. 207.

Cones whitish, pendent. Mond. Poly.

Scotch Fir. Highland mountains.

FUMA'RIA. See Gen. Ch. p. 60.

F. Seed-vessels in bunches, 1 seed officinalis. in each: stem spreading.

With. Arr. 620. Curt. 2.

Vith. Arr. 620. Curt. 2. Woodv. 88.

Flowers pink and dark purple.

Common Fumitory.

Aug. Diad. Hex.

Corn-fields, hedge-banks.

Polyg'Ala. Cal. 5-leaved, 2 larger wing-like, before the ripening of the feeds, coloured: Capf. inverfely heart-fhaped, 2-celled: feeds folitary.

vulgáris.

P. Flowers in bunches: stems herbaceous, simple, trailing: leaves strap-spear-shaped.

With. Arr. 622. E. Bot. 76.

Bloff. blue, purple, flesh colour, or white.

Diad. Oct.

Common Milkwort. Paftures, heaths. July.

SPAR'TIUM. Cal. extending downwards, 2-lipped: Filaments adhering to the germen: Summit woolly above.

fcopárium.

S. Leaves in threes, folitary: branches without prickles, angular.

With. Arr. 623. Curt. 5. Woodv. 89.

Diad. Dec.

Bloff. yellow. June. Common Broom. Dry pastures.

GENISTA. Cal. 2-lipped, the upper 2, the lower 3-toothed: Standard oblong, bent back from the stamens and pistil: Legumen regularly-shaped.

G. Leaves spear-shaped, blunt: pilosastem with tubercles, prostrate. With. Arr. 625. E. Bot. 208.

Bloff. yellow. Diad. Dec. Paftures, heaths. June.

G. Very thorny; fmooth: thorns anglicafimple and compound: leaves fpear-shaped.

With, Arr. 625. E. Bot. 132.

Flowers pale yellow. June.

Needle Furze. Petty Whin. Diad. Dec.

Heaths, moist spongy ground.

ANTHYL'LIS. Cal. bellying: Legumen roundish, covered.

A. Herbaceous: leaves winged, Vulnerária. unequal: flowers in a double head.

With. Arr. 629. E. Bot. 104.

Flowers yellow.

Kidney Vetch. Ladies Finger.

Meadows, pastures, in a calcareous foil. Aug. Diad. Dec.

Var. 2. Bloffoms fcarlet.

O'ROBUS. See Gen. Ch. p. 61.

fylvat'icus. O. Stems drooping, hairy, branched: leafits 7 to 12 pair.

With. Arr. 630.

Bloff. long and narrow. July.

Diad. Dec. Bitter Vetch. Wood Peafeling. Woods, hedges, pastures.

LATHYRUS. Cal. 2 upper fegments fhorter: Style flat, broader upwards, woolly on the upper furface: Legumen generally equal, broad.

Niffolia. L. Leaves fimple: flipfulæ awlfhaped.

With. Arr. 632. E. Bot. 112.

Bloff. beautiful crimfon.

Diad. Dec. Crimfon Grafs Vetch. Borders of corn-fields, pastures.

Vícia. See Gen. Ch. p. 61.

fépium. V. Legumens mostly 4 together, upright: leasits egg-shaped, very entire, the outer ones gradually smaller.

With. Arr. 637.

Bloff. dirty purple. June. Diad. Dec. Bush Vetch. Woods, hedges.

V. Legumens folitary, upright, lathyroides. fmooth; leafits about 6, the lower ones inverfely heart shaped: stipulæ half arrow-shaped, very entire: seeds cubic, warty.

With. Arr. 638. E. Bot. 30.

Flowers fmall, purplish blue.

Strangle Vetch, or Tare. Dry Diad. Dec. pastures, gravel pits.

ORNITHOPUS. Seed-veff. cylindrical, but jointed and crooked.

O. Leaves winged: legumens bow- perpufil'lus. ed in, much larger than the winged flower-scale.

With. Arr. 640. Curt. 6. E. Bot. 369.

Common Birds'-foot. Sept. Diad. Dec. Sandy banks, road fides.

HIPPOCRE'PIS. Seed-veffel many-celled, compressed, crooked, with

feveral notches along one of the feams.

eomósa. H. Legumens on fruit - stalks, crowded, bowed, serpentine.

With. Arr. 641. E. Bot. 31.

Bloff. dull yellow.

Dia. Dec. Tufted Horse-shoe Vetch. Meadows in a calcareous foil.

HEDYS'ARUM. Keel broad and blunt on the outer part: Legumen jointed, 1 feed in each joint.

Onob'rychis H. Leaves winged: legumens with 1 feed, prickly: wings as long as the calyx: stem growing long.

With. Arr. 642. E. Bot. 96.

Diad. Dec. Common Saintfoin. Cock's-head.

Meadows in chalky foil. July.

TRIFO'LIUM. Flowers mostly forming a head: Capf. or Legumen, scarcely longer than the calyx; not opening, but falling off entire. T. Heads like umbels: legumens répens.

4-seeded: stem creeping.

With Arr. 646. Curt. 3.

Bloff. white. Sept. Diad. Dec.

White Trefoil. Dutch Clover.

T. Heads like umbels: legumens by bridum.

4-feeded: ftem afcending.

With. Arr. 647.

Moift pastures. Sept. Diad. Dec.

T. Heads woolly: 5-flowered; a fubterrárigid tuft from the centre reflected neum. upon, and enclosing the fruit. With. Arr. 647. Curt. 2.

Bloff. white.

Subterraneous Trefoil. Aug. Diad. Dec. Heaths in fandy gravelly foil.

T. Heads fitting, hemispherical, glomerátum, rigid: calyxes scored, teeth expanded equal.

With. Arr. 648. Curt. 4.

Bloff. pale red. June. Diad. Dec.

Round - headed Trefoil. Sandy meadows, heaths.

MAY.

fcábrum. T. Heads fitting, lateral eggfhaped: fegments of the calyx unequal, stiff, bowed back. With. Arr. 648. Curt. 6.

Bloff. whitish.

Diad. Dec. Hard-knotted Trefoil. June. Chalky, fandy pastures.

praten fe. T. Spikes crowded: blossoms unequal: calyx with 4 of the teeth equal: stipulæ awned: stems ascending.

With. Arr. 651.

Bloff. reddish purple, or white.

Diad. Dec. Purple Trefoil. Honeyfuckle Trefoil. Meadows, pastures. Sept.

filifor me. T. Spikes oval, loofely tiled, fewflowered: flems trailing.

With. Arr. 655.

Bloff pale yellow.

Diad. Dec. Small Trefoil. Hop Trefoil. Sandy heaths, pastures. July.

Medica'Go. Piftil bent, preffing down the keel and springing out of it with a jerk. Seed-veffel a legumen, compressed, bent, or twisted spirally.

M. Spikes oval: feed-veffels kid- lupulina.
ney-shaped, with 1 cell and 1
feed: stems trailing.

With. Arr. 659. Curt. 2.

Bloff. yellow. Aug. Diad. Dec. Trefoil Medick. Melilot Trefoil. Black Nonfuch. Norfolk. Cornfields, meadows, pastures.

M. Fruit-stalks with 2 or 3 seed- arábica. vessels: legumens prickly: stipulæ toothed: leasits inversely heart-shaped: stems spreading.

With. Arr. 660. Curt. 3.

Bloff. yellow.

Heart Trefoil. Heart Clover. Diad. Dec. Dry fandy pastures. June.

M. Legumens mostly three, on a min'ima. fruit-stalk: prickles hooked: stipulæ entire.

With. Arr. 660.

Bloff. fmall, pale yellow. Diad. Dec. Sandy places.

muricáta. M. Fruit-stalks many - flowered: legumens roundish, thorny, hoary: leaves woolly.

Diad. Dec. With. Arr. 664.

At Orford in Suffolk on the fea

bank.

TRAGOPO GON. Recept. naked: Calyx fimple: Down feathered.

porrifólium. T. Calyx longer than the rays of the bloffom: leaves entire, ftiff and ftraight: fruit-ftalks thickening upwards: florets very narrow, lopped.

With. Arr. 672.

Bloff. purple.

Syng. Æqua. Purple Goats'-beard. Salfafy. Meadows, pastures.

LEON TODON. See Gen. Ch. p. g.

his pidum. L. All the scales of the calyx upright: leaves toothed, very entire, hispid: bristle-like hairs forked.

With. Arr. 681. Curt. 5.

Syng. Æqua. Rough Dandelion.

Meadows, paftures. Sept.

HIERA'CIUM. Recept. generally naked: Calyx tiled, egg-shaped, fometimes double: Down mostly sitting, hair-like: rarely feathered.

H. Leaves very entire, egg-fhap- pilofel'la. ed, cottony underneath: fuckers creeping.

With. Arr. 682. Curt. 4.

Bloff. pale yellow within.

Creeping Moufe-ear. Dry Mea- Syng. Æqua. dows, walls. Sept.

Hyo'seris. Recept. naked: Calyx nearly equal: Down fitting, like awned chaff, furrounded with hairs; or, instead thereof, a minute calyx including a very fine down.

H. Stem divided, naked: fruit-min'ima. stalks thickening upwards.

With. Arr. 690. E. Bot. 95.

Bloff. yellow.

Small Swine's Succory. June. Syng. Æqua.

Meadows in gravelly foil.

HYPOCHERIS. Recept. chaffy: Cal. somewhat tiled: Down on a pedicle; feathered.

radicáta. H. Stem branched, naked, even; leaves notched, blunt, rough; fruit-stalks scaly.

With. Arr. 692. Curt. 3.

Bloff. yellow within, reddishgreen without.

Syng. Æqua. Long-rooted Hawkweed Meadows, pastures. cpt.

CAR DUUS. Cal. bellying illed : feales thorny: Recept. hairy.

praten fis. C. Leaves spear-shaped, ragged, and edged with unequal prickles: cottony underneath: stem cottony, generally with 1 slower and 2 leaves.

With. Arr. 701. E. Bot. 177.

D

Bloff. red. June.

Syng Equa. Meadow Thiftle. English Soft, or Gentle Thistle. Meadows.

GNAPHA'LIUM. Recept. naked: Down hair-like, or feathered: Calyx tiled: scales at the edge roundish, skinny, coloured.

G. Runners trailing: stem undivided: flowers in a simple corymbus: male and semale flowers on distinct plants.
Male plant.

With. Arr. 712. E. Bot. 267.

Bloff. white, purple, or red.

Cats'-foot. Cudweed. June. Syng. Super.

Dry mountainous pastures.

CINERA'RIA. Recept. naked: Down hair-like: Cal. fingle, many-leaved, equal.

C. Flowers in a terminating um-integrifdia.
bel, with an involucrum at its
base: leaves inversely egg-shaped, woolly, indistinctly toothed.

With. Arr. 730. E. Bot. 152.

Flowers 1 to 4, yellow.

-

d

7.

e.

or

d:

d:

Mountain Fleawort. Hilly paf- Syng. Super. tures. June.

Doro'nicum. Recept. naked: Down hair-like: Cal. scales in 2 rows; equal, longer than the blossoms:

L 2

Seeds of the circumference downless and naked.

Pardalian'- D. Leaves heart - shaped, blunt, finely toothed: root-leaves on leaf-stalks: stem-leaves embracing the stem.

With. Arr. 732.

Syng, Super. Bloff. yellow. June.

Great Leopard's Bane. Wolf's
Bane.

MATRICA'RIA. Recept. naked: Down none: Calyx hemispherical, tiled; fcales rather pointed, not skinny at the edge.

Chamomil'la. M. Receptacles conical; rays expanding: calyx scales equal at the edge.

With. Arr. 736. Curt. 5.

Syng. Super.

Chamomile Feverfew. Aug. Corn-fields, dunghills, road-fides.

AN'THEMIS. Recept. chaffy: Down none: Calyx hemispherical, scales nearly equal: Florets of the circumference more than 5.

1016 ,014

A. Receptacles conical: chaff brif- Cótula. tle-shaped: feeds naked.

With. Arr. 738. Curt. 5.

Stinking Chamomile. Mayweed.

Corn-fields, road-fides, dunghills.

book sale to an analytic

AD VIE WINGSTON

Zawarnet Line Flowers Meleym I Few on the time plants without

the state of the s

District of the state of

Var. 2. Double-flowered. All the Syng. Super: florets narrow. Aug.

JUNE.

CHA'RA. Male. Calyx none: Anther at the base of the germen. Fem. Cal. 4-leaved: Summits 5cleft: Seed 1.

omentósa. C. Prickles on the stem eggshaped.

With. Arr. 2. Oct.

Mo. Mo. Brittle Storewort. Ditches, ponds.

his pida. C. Prickles on the stem hair-like, in clusters.

With. Arr. 3. Oct.

Mo. Mo. Prickly Stonewort. Ditches, pools.

fléxilis.

C. Joints of the stem without prickles, semi-transparent, broader upwards. Of an equal thickness the whole length.

With. Arr. 3. Oct.

Mo. Mo. Smooth Stonewort. Ponds, ditches.

ZANNICHEL'LIA. Flowers Male and Fem. on the fame plant: without petals.

Male. Cal. none.

Fem. Cal. 1 leaf, bell-shaped:

Z.

palústris.

With. Arr. 6.

Horned Lakeweed, or Pondweed.

Ditches, stagnant waters. July. Mo. Tet.

With 2 cells, 2 feeds in each cell.

L. Leaves egg-fhaped, blunt: pa- vulgáre. nicle with 3 divisions.

With. Arr. 10. Curt. 5.

Bloff. white. Berries black.

Privet. Prim. Print. Hedges in Di. Mo. gravelly foil.

CIRCE A. Bloff. 2 petals: Cal. 2leaved, fuperior.

C. Stem upright: bunches feve- lutetiána. ral: leaves egg-shaped, somewhat serrated: rough with hair. Leaves egg-spear-shaped, hairs somewhat serrated.

With, Arr. 10. Curt. 3.

Di. Mo.

spicata.

JUNE.

Enchanter's Nightshade. July. Woods, moift hedge bottoms.

VERO'NICA. See Gen. Ch. p. 5.

Spike terminating: leaves opposite, blunt, scolloped: stem ascending, undivided: spike conical: leaves fpear-shaped: stamens much longer than the bloffoms.

With, Arr. 12. E. Bot. 2.

Bloff. blue.

Upright-spiked Male Speedwell. Di. Mo. Mountainous moist pastures.

> V. Bunch terminating, fomewhat like a corymbus: leaves oppofite, nearly egg - shaped, smoothish, flightly fcolloped: fruit - ftalks longer than the floral-leaves: ftems and cups woolly.

> > With. Arr. 14.

Bloff. blue-red with purple lines. Di. Mo. Rocks. July.

Bunches lateral: leaves egg-Beccabun'ga. V. shaped, flat: stem creeping. With. Arr. 15. Curt. 2. Woodv. 7.

faxátilis.

Bloff. blue.

Brooklime. Slow shallow streams. Di. Mo.

V. Bunches lateral, thinly fet with fcutelláta. flowers, wide-fpreading, alternate: little fruit-stalks pendent: leaves very entire, strap-shaped.

With. Arr. 16. Curt. 5.

Bloff. white or purplish.

Narrow-leaved Speedwell. Poor Di. Mo. fwampy foil. Aug.

PINGUI CULA. Bloff. gaping, ending in a fpur: Cal. 2-lipped, 5-cleft: Capf. 1 cell.

P. Nectary blunt, shorter than the lusitan'icapetal: stalk hairy: capsule globular: nectary thickest at the end.

With. Arr. 18. E. Bot. 145.

Bloff. lilac-coloured.

Marthes.

stalk.

Di, Mo.

July.

SAL'VIA. Bloff. gaping: Filaments fixed transversely to a little foot-

Di. Mo.

praten is. S. Leaves heart-oblong, scolloped:
upper leaves embracing the stem:
flowers in whirls with hardly any
intermixture of leaves: helmet of
the blossom gummy.

With. Arr. 21. E. Bot. 153.

Bloff. bluish purple.

Di. Mo. Meadow Sage, or Clary. Meadows, pastures. July.

verbenáca. S. Leaves indented, ferrated, fmoothish: blossom more slender than the calyx.

With. Arr. 21. Curt. 6. E. Bot. 154.

Bloff. blue.
Wild Clary. Wild Sage.

OR'CHIS. See Gen. Ch. p. 30.

bifólia. O. Lip of the nectary spear-shaped, very entire: horn very long: petals expanding.

With. Arr. 21. E. Bot. 22.

Flowers white, of a sweet scent in the night.

Di. Mo. Butterfly Orchis. Meadows, at Sedgwick Park, Suffex.

O. Lip of the nectary 2-horned, pyramidális.

3-cleft: fegments equal, very entire: horn long: petals fomewhat spear-shaped.

With. Arr. 23. E. Bot. 110.

Flowers purple.

Late-flowering Orchis. July. Di. Mo.

Meadows in a calcareous foil.

O. Bulbs expanding: horn of the maculata. nectary shorter than the germen: lip flat: petals on the back upright.

With. Arr. 27.

Flowers pale purple.

Female-handed Orchis. Spotted Orchis.

Di. Mo.

Woods, rich meadows.

O. Horn of the nectary briftle- conop'sea. fhaped, longer than the germen: lip cloven into 3: two of the petals very much expanding.

With. Arr. 28. E. Bot. 10.

Flowers flesh - coloured, or pale purple, very sweet.

Red-handed Orchis. Meadows. Di. Mo.

SATY'RIUM. Nectary behind the flower: inflated, roundish.

hircínum. S. Bulbs undivided: leaves spearshaped: lip of the nectary 3-clest: the middle segment strap-shaped, oblique, bitten off.

With. Arr. 29. E. Bot. 34.

Flowers whitish, smelling like a goat.

Di. Mo. Lizard Flower, or Satyrion.

Chalky meadows. July.

viride. S. Bulbs hand-shaped: leaves oblong, blunt: lip of the nectary strap-shaped, 3-cleft: the middle segment indistinct.

With. Arr. 30. E. Bot. 94.

Flowers pale, greenish.

Di. Mo. Frog Satyrion. Aug.

Meadows, in gravelly foil.

fhaped: lip of the nectary 3-cleft: fegments acute: the middle one blunt.

Di. Mo.

With, Arr. 31.

White Satyrion. July. Di. Mo.

Moift meadows in Wales.

S. Roots fibrous: leaves egg-shap- repens. ed, growing from the root: flowers pointing one way.

With. Arr. 13. E. Bot. 289.

Flowers whitish, or straw-colour.

Creeping Satyrion. Old mosfy Di. Mo. Aug. woods.

O'PHRYS. See Gen. Ch. p. 34.

O. Bulb fibrous: ftem with 2 cordáta. leaves: leaves heart-shaped.

With. Arr. 35. E. Bot. 358.

Leaft Twayblade. July. Di. Mo.

Moift mountainous heaths.

O. Bulb roundish: stalk naked, Loeselii. 3-cornered: lip of the nectary egg-fhaped.

With. Arr. 35. E. Bot. 47.

Flowers in a bunch, yellowish July. Di. Mo. green.

Dwarf Twayblade. Moors.

O. Bulbs roundish: stalk leafy: anthropophóra. lips of the nectary strap-shaped,

3-cleft: the middle fegment long, cloven.

With. Arr. 37. E. Bot. 29.

Di. Mo. Man Twayblade. Chalky meadows.

apiféra. O. Bulbs roundish: stem leafy:
lip of the nectary with lobes, the
lobes bent in underneath, shorter
than the petals.

With. Arr. 38. Curt. 1. E. Bot. 383.

Di. Mo. Bee Orchis, or Twayblade. Meadows.

SERA PIAS. See Gen. Ch. p. 71.

longifolia. S. Leaves fpear - fhaped, embracing the stem: flowers nearly horizontal: lip scolloped, blunt, longer than the petals.

With. Arr. 41. E. Bot. 270.

Marsh Helleborine. Aug.

Di. Mo. Marthy places.

MALAX'IS. Nectary 1 leaf, heartfhaped, concave, enclosing the stamens and pistils. M. Stalk 5-edged: leaves fheath- paludofa-

With. Arr. 39. E. Bot. 72.

Flowers green yellow.

Marsh Twayblade. Near Nor- Di. Mo. wich. July.

LEM'NA. See Gen. Ch. p. 71.

L. Leaves on leaf-stalks, spear- triful ca. shaped.

With. Arr. 43.

Ivy-leaved Duckmeat. Sept.

Ditches, flagnant waters. Di. Mo.

L. Leaves fitting, flattish on both minor. fides: roots solitary.

With. Arr. 44.

Least Duck-meat. Greeds. Pools. Di. Mo. Flowers in the dog-days.

Sept.

Sa'LIX. See Gen. Ch. p. 31.

S. Leaves egg - shaped: semi- myrsinites. transparent.

With. Arr. 48.

Whortle-leaved Willow. High- Di. Mo. land mountains.

JUNE.

retúsa.

S. Leaves fomewhat ferrated, inverfely egg-shaped, very blunt.

With. Arr. 48.

Di. Mo.

Rocks in Scotland.

lappónum.

S. Leaves hairy, spear-shaped.

With. Arr. 51.

Di. Mo.

Lapland Willow. Highland mountains.

VALERIA'NA. See Gen. Ch. p. 34.

officinális.

V. Leaves all winged and toothed. With. Arr. 66. Curt. 6. Woodv. 96.

Bloff. pink.

Tri. Mo.

Great Wild Valerian. Hedges, woods, marshes.

I'ms. Bloff. with 6 divisions, unequal, every other segment bent back as if jointed: Summits petallike, 2-lipped, edges at the base turned up.

fœ tida.

I. Stem with 1 angle.

With. Arr. 70.

Bloff. purple ash-colour.

Var. 2. Leaves striped.

Stinking Gladdon. Hedge-banks. Tri. Mo. July.

I. Every other fegment of the psuedácorus. blossom smaller than the summit.

With. Arr. 69. Curt. 3.

Woodv. 40.

Bloff. yellow.

Flower-de-luce. Water Flag. Tri. Mo. Banks of rivers. July.

NAR DUS. Cal. none: Bloff. 2 valves: Nectary none.

N. Spike slender, straight: the stric'ta. slorets pointing in one direction.

With. Arr. 71. E. Bot. 290.

Small Matweed. Heaths, marshes. Tri. Mo.

ERIO'PHORUM. See Gen. Ch. p. 10.

E. Straws cylindrical: leaves channelled and 3-cornered: fpikes on fruit-stalks.

With. Arr. 72.

Cotton-grass. Moor-grass. Tri. Mo. Marshes, bogs.

M

alpinum.

E. Straws naked, 3-cornered: spike shorter than the down.

With. Arr. 72. E. Bot. 311.

Moffes.

DIP'SACUS. Cal. common, manyleaved: proper, superior: Recept. chaffy.

sylvéstris.

D. Leaves in opposite pairs, united at the base: chaff straight.

With. Arr. 182. Curt. 3.

Tet. Mo.

Wild Teafel. Wet hedge-banks.

Aug.

fullónum.

D. Leaves fitting: ferrated: chaff bent backwards.

With. Arr. 182.

Tet. Mb.

Manured Teafel.

Scabio'sa. Cal. common, manyleaved: proper, double, superior: Recept. naked, or chaffy: Seed wrapped in the proper cup.

fuccis'a.

S. Bloffoms 4-cleft, equal: ftem undivided: branches approaching: leaves spear-egg-shaped.

With. Arr. 183. Curt. 3.

Bloff. bluish purple.

Devil's-bit Scabious.
Fields, pastures.

Aug. Tet. Mo.

S. Bloffoms 5-cleft, radiating: columbária. root-leaves fimple, fcolloped: ftem-leaves compound.

With. Arr. 184.

Flowers lilac-colour.

Small Scabions.

Sept. Tet. Mo.

Dry hilly pastures.

S. Bloffoms 4-cleft, radiating: arven'fis. leaves wing-cleft and jagged: ftem rough with strong hairs.

With. Arr. 183. Curt. 4.

Bloff. blue, purple, or white.

Field Scabious. Pastures, corn- Tet. Mo. fields. Aug.

ASPE'RULA. See Gen. Ch. p. 77.

A. Leaves 4 in a whirl, flrap-cynan'chica. fhaped: upper leaves opposite: flem upright: flowers 4-cleft.

With. Arr. 186. E. Bot. 33.

Bloff. pale rose red.

M 2

Squinancy. Wort. Tet. Mo. July. Limestone hills, chalky foil.

GA'LIUM. See Gen. Ch. p. 77.

erectúm. G. Leaves mostly 8 in a whirl: fpear-shaped, with fine prickly ferratures: panicles with 3 divisions: tem rough, flaccid.

With. Arr. 189.

Flowers white, 4-cleft.

Tet. Mo. Upright Goofe-grafs. Meadows, wet pastures. July.

mollúgo. G. Leaves 8 in a whirl, egg-ftrapfhaped, dagger-pointed: fomewhat ferrated, greatly expanded: stem feeble: branches greatly expanding.

With. Arr. 189.

Bloff. white.

Tet. Mo. Madder Goofe-grafs. White Ladies Bedftraw. Hedges, heaths. Aug.

G. Male flowers 3-cleft, on pespurium. dicles fpringing from the fruitstalks of the hermaphrodite flowers.

With. Arr. 190. July.

Corn-fields, in a limestone soil. Tet. Mo.

G. Leaves about 6 in a whirl, anglicum. fpear-shaped, pointed, edged with prickles, reflected: stem wide-spreading, rough with prickles pointing downwards.

With. Arr. 191. E. Bot. 384. Tet. Mo.

Bloff. greenish yellow.

Small Goofe-grafs. Walls. July.

G. Leaves unequal, strap-spear-palus tre. shaped, smooth, entire, blunt, 4, 5, or 6, in a whirl: stems rough, spreading, branched upwards.

With Arr. 187.

White Ladies Bedstraw. July. Tet. Mo. Banks of rivulets and pools.

G. Leaves 8 in a whirl, ftrap-shap-vérum. ed, furrowed: flowering branches short.

With Arr. 191. Curt. 5.

Yellow Ladies Bedstraw. Cheese- Tet. Mo. rening. Sides of fields and roads.

LITTOREL'LA. M. and Fem. flowers on the same plant.

Male. Bloff. 4-cleft: Cal. 4 leaved: Stam. long.

Fem. Bloff. mostly 4-cleft: Cal. none: Style long: Seed a nut.

lacus tris. L.

With. Arr. 194.

Tet. Mo. Plantain Shoreweed. Marshy places, in fandy foil.

PLANTA'GO. Bloff. 4-cleft, permanent, the border broken back: Cal. 4-cleft: Stam. extremely long: Capf. 2-celled, cut round, fuperior.

májor.

P. Leaves egg-shaped, smooth: stalk cylindrical: spike tiled with florets.

With. Arr. 195. Curt. 2. Woodv. 14.

E

Tet. Mo.

Way-bread. Great Plantain.
Road-fides. Aug.

média. P. Leaves egg-spear-shaped: pu-

bescent: spike and stalk cylindrical.

With. Arr. 196. Curt. 4.

Lamb's Lettuce. Hoary Plantain.
Road-fides; pastures, in chalky, Tet. Mo.
marly soils. Aug.

P. Leaves spear - shaped: spike lanceoláta. nearly egg-shaped, naked: stalk angular.

With. Arr. 196. Curt. 2.

Ribwort Plantain. Pastures. Tet. Mo.

P. Leaves strap-shaped, mostly entire: woolly at the base: spike and stalk cylindrical.

With. Arr. 197. E. Bot. 175. Tet. Mo. Sea Plantain. Sea coast. July.

P. Leaves strap shaped, toothed; coron opus. stalk cylindrical. Leaves between winged and toothed.

With. Arr. 198.

Star of the Earth. Hartshorn. Tet. Mo. Gravelly soil, sea shore. Aug.

EPIME DIUM. Bloff. 4 petals: Nectaries 4, leaning: Cal. deciduous: Seed-veffel a pod. alpinum.

E.

With. Arr. 199.

Tet Mo. Bloff. mulberry red with a yellow

Bingley woods.

July.

CENTUN'CULUS. Cal. 4-cleft: Bloff. 4-cleft, expanding: Stam. short: Capf. 1-celled, cut round.

min imus.

Tet. Mo.

C.

With. Arr. 198 Curt. 3.

Bloff. minute, white.

Bastard Pimpernel. Moist fandy ground.

SANGUISOR BA. Bloff. 4-cleft: Germen betwixt the calyx and the bloffom.

officinalis.

S. Spikes egg-shaped.

With. Arr. 199.

Tet. Mo.

Wild, or Great Burnet. Aug. Pastures in calcareous soil.

COR'NUS. Involucrum generally 4leaved: Petals 4, superior: Drupa fucculent, beneath, 2-celled, hard, solitary. C. Branches straight, leaves egg- fanguin'eashaped, green on both sides: tuft flatted.

With. Arr. 200. E. Bot. 249.

Bloff. white.

Tet. Mo.

Dogberry Tree. Woods, hedges.

C. Herbaceous, with 1 pair of suécia. branches.

With. Arr. 200. E. Bot. 310.

Bloff. white.

Dwarf Honeysuckle. Hills in the Tet. Mo. North. July.

ALCHEMIL'LA. Cal. 8-cleft: Bloff. none: Seed 1 or 2 enclosed by the calyx.

A. Leaves gashed.

vulgáris.

With. Arr. 204.

Flowers yellowish green.

Bearsfoot. Common Ladies Mantle. Tet. Mo. Meadows, pastures. Sept.

POTAMOGE'TON. See Gen. Ch. p. 80.

perfoliatum. P. Leaves heart-shaped, embracing the stem.

With. Arr. 212. E. Bot. 168.

Bloff. yellowish.

Tet. Tet. Perfoliate Pondweed. Slow rivers, ponds. Aug.

lúcens. P. Leaves very long and spearshaped, fitting: upper ones opposite.

With. Arr. 212. E. Bot. 376.

Bloff. reddish.

Tet. Tet. Long-leaved Pondweed. Rivers, ponds. July.

compress - P. Leaves strap-shaped, blunt: stem flatted.

With. Arr. 213.

Bloff. greenish.

Tet. Tet. Flat-flalked Pondweed. Slow ftreams, ditches. July.

pusiflum. P. Leaves strap-shaped, opposite and alternate, distinct, expanding from the base: stem cylindrical.

With. Arr. 214. E. Bot. 215.

Tet. Tet. Small Pondweed. Ponds in clayey foil. Aug.

SAGI'NA. See Gen. Ch. p. 37.

S. Branches trailing. procum bens With. Arr. 215. Curt. 3.

Bloff. greenish white.

Chickweed. Breakstone. Walls, Tet. Tet. roofs, fand and boggy places.

S. Stem spreading, forked: leaves cerastoides. battledore, or inversely egg-shaped: pointed, bent back: fruitstalks reflected after flowering.

With. Arr. 216. E. Bot. 166.

Bloff. 4 petals, white. Tet. Tet.

Cynoglos'sum. See Gen. Ch. p. 84.

C. Stamens shorter than the blof-officinale. fom: leaves broad, spear-shaped, fitting, cottony.

With. Arr. 227. Curt. 4. Woodv. 216.

Bloff.marone, or mulberry-colour.

Great Houndstongue. Road fides, Pen. Mo. rubbish.

Vi ola. See Gen. Ch. p. 6.

*palus tris. V. Leaves kidney-shaped.

With. Arr. 261. Curt. 3.

Pen. Mo. Marsh Violet. Moist meadows, bogs. April.

Bora'go. Bloff. wheel - shaped; mouth closed with rays: Nuts 4, not perforated.

officinális. B. All the leaves alternate: calyxes expanding.

With. Arr. 230. Woodv. 217.

E. Bot. 36.

Pen. Mo. Common Borage. Walls, rubbish. Aug.

Lyco'rsis. Bloff. tube crooked:

Mouth closed with scales: Nuts
4, perforated.

arven fis. L. Leaves spear-shaped, rough with hair: calyx while in flower upright.

With. Arr. 231. Curt. 5.

Bloff. fky-blue.

Pen. Mo. Wild Bugloss. Corn-fields, roadfides. Sept.

* This plant was by mistake omitted in March, when it first flowers. It should have followed Violo hirta, p. 18.

E'CHIUM. Bloff. irregular: Mouth naked: Nuts 4, large, rough, imperforated.

E. Stem rough with briftles and vulgare. tubercles: ftem - leaves spearshaped, rough with hair: flowers in lateral spikes. With. Arr. 232. E. Bot. 181.

Bloff. before it expands bright red, afterward blue.

Common Viper-grass. Sandy corn- Pen. Mo. fields, walls, rubbish. Aug.

E. Stem upright, hairy: fpikes ital'icum. hairy: fegments of the bloffoms nearly equal: stamens extremely long.

With, Arr. 232.

Bloff. white.

5

1

.

n

2

Wall Viper-grafs. Sandy meadows, Pen. Mo. pastures.

Bloff. hairy, or MENYAN THES. fringed: Nect. 5 at the base of the germen: Summit 2-lobed: Capf. 1-celled.

JUNE.

trifoliáta.

M. Leaves growing by threes: bloffom fegments entire at the edge; shaggy on the upper furface.

With. Arr. 236. Curt. 4. Woodv. 2.

Bloff. pinky, and white.

Pen. Mo.

Marsh Trefoil. Ponds, pits.

July.

C

C

Z

Stem fixed to the top of the tube: Summit globular: Capf. 1-celled, globular.

palus tris.

H. Fruit-stalks in whirls, many-flowered.

With. Arr. 236. Curt. 1. E. Bot. 364.

Pen. Mo.

Water Violet. Water Yarrow.
Ponds, ditches. July.

Lysima Chia. Bloff. wheel-shaped: Caps. globular; dagger-pointed: of 1-celled, 5 valves, and many seeds.

vulgáris.

L. Panicled: bunches terminating. With, Arr. 237. Curt. 5. Flowers yellow.

Yellow Willow-herh, or Loofefirife. Banks of rivers. July. Pen. Mo.

L. Bunches lateral, on fruit- thyrfiflóra. ftalks.

With. Arr. 237. E. Bot. 176.

Bloff. yellow.

Tufted Loofestrife. Marshes, Pen. Mo. banks of rivers.

L. Leaves fomewhat heart-shaped: Nummuláflowers folitary: stem creeping. ria.

With. Arr. 238. Curt. 3.

Bloff. yellow.

Money-wort. Herb Twopence. Pen. Mo. Meadows.

Convol'vulus. Bloff. bell-shaped, plaited: Nect. surrounding the base of the germen: Summits 2: Caps. 2 or 3-celled, 2 seeds in each.

C. Leaves arrow fhaped, acute on arven fis. each fide: fruit-stalks bearing generally 1 flower.

With. Arr. 239. Curt. 2. E. Bot. 312.

Bloff. flesh-coloured, almost white.

Pen. Mo., Small Bindweed. Corn-fields, road-fides. July.

POLEMO'NIUM. Bloff. wheel-shaped, with 5 divisions: Filaments broad and membranaceous at the base: Summit 3-cleft: Caps. 3-celled, opening at the top: Seeds angular.

right: calyx longer than the tube of the bloffom.

With. Arr. 241. E. Bot. 14.

Bloff. blue, or white.

Pen. Mo. Greek Valerian. Jacob's Ladder.

CAMPA'NULA. Bloff. bell-shaped: Filaments broad and arched at the base: Summit 3-cleft: Caps. beneath, 3-celled, opening at 3 lateral holes.

what branched at the base: leaves oblong, scolloped: cups incorporated, and longer than the blossom.

With. Arr. 244.

Bloff. purple.

Leffer Venus' Looking-glafs.

Chalky corn-fields. July. Pen. Mo.

Sa'molus. Bloff. falver - shaped: Stamens protected by the valves of the blossom: Capf. 1-celled: beneath: opening with 5 valves at the top.

S. With. Arr. 246. Curt. 4.

Bloff. white.

Round-leaved Water Pimpernel.

Marshes, moist meadows. July. Pen. Mo.

JASIO'NE. Common Calyx 10-leaved: Cup proper, 5 toothed: Bloff. 5 petals regular: Anthers united: Capf. beneath; 2-celled; manyfeeded, opening at the top, crowned by the proper cup.

J. Leaves strap shaped, very entire. montána. With. Arr. 247. Curt. 4.

Bloff. blue, or white.

Hairy Sheep Scabious. Sandy, bar- Pen. Mo. ren fallows. July:

Pen. Mo.

VERBAS CUM. Bloff. wheel-shaped, nearly regular: Capf. 2-celled, 2-valved, many-seeded.

Thap'sus. V. Leaves decurrent, cottony on both sides: stem unbranched: fummit globular.

With. Arr. 248. Woodv. 125.

Bloff. yellow, rarely white.

Great White Mullein. Dry ditchbanks, in chalky foil. July.

Blattária. V. Leaves embracing the stem, oblong, smooth: fruit-stalks solitary.

With. Arr. 251.

Pen. Mo. Yellow Moth Mullein. Gravelly foil. July.

HYOSCY'AMUS. Bloff. funnel-shaped, blunt, irregular: Stam. leaning: Capf. with a lid, 2-celled: Seeds many, kidney-shaped.

niger. H. Leaves embracing the stem, indented: flowers sitting.

With. Arr. 252. Woodv. 52.

S

Pen. Mo. Common Henbane. Road-fides, rubbish.

ATROPA. Bloff. bell-shaped: Stam. distant: Berry globular, 2-celled.

A. Stem herbaceous: leaves egg- Belladon'na. shaped, entire.

With. Arr. 252. Curt. 5.

Bloff. dark purple.

1.

2. es, Deadly, or Sleepy Nightshade.

Hedges, limestone rubbish.

Pen. Mo.

Aug.

Sola'Num. Bloff. wheel-shaped:

Anthers a little united, 2 holes
at the top of each: Berry 2celled.

S. Stem without prickles, rather Dulcamára. fhrub-like, zigzag: upper leaves halberd-fhaped: flowers in tuft-like bunches.

With. Arr. 253. Curt. 1. Woodv. 33.

Bloff. purple. Berries scarlet.

Bitter-sweet. Woody Nightshade. Pen. Mo.
Hedges, fides of ditches. July.

S. Stem without prickles, her-nigrum. baceous: leaves egg - shaped, N 2 Fen. Mo:

toothed, angular: bunches nodding, pointing 2 ways.

> With. Arr. 254. Curt. 2. Woodv. 226.

Bloff. white. Berries black.

Common, or Garden Night shade.

Rubbish, dunghills, gardens.

Oct.

CHIRO'NIA. Bloff. funnel-shaped: Piftil leaning: Stam fixed on the top of the tube of the bloffom: Anthers becoming spiral: Seedveffel 2-celled.

Centaúrium C. Herbaceous: leaves spear-shap ed: calyx fhorter than the tube of the bloffom.

> With. Arr. 255. Curt. 4. Woodv. 157.

> > I

Bloff. yellowish, border pinky red.

Leffer Centory. Barren pastures. Pen. Mo. Aug.

> GLAU'X. Cal. 1 leaf: Bloff. none: Capf. 1-celled, 5-valved, 5-feeded.

G.

marit'ima.

With. Arr. 268. E. Bot. 13.

Flowers numerous, purplish, or whitish.

Sea Milkwort. Black Saltwort.
Salt marshes. July. Pen. Mo.

THE SIUM. Cal. 1 leaf, on which ftamens grow: Seed 1 beneath.

T. Spike branched: floral leaves linophyl'by threes: leaves ftrap-spearshaped: calyx tube very short. With. Arr. 268. E. Bot. 247.

Dry heaths and hills in a calcareous foil. July. Pen. Mo.

Hu'mulus. Male and Fem. flowers on different plants: Bloff.

Male. Cal. 5-leaved.

Fem. Cal. 1-leaved, with a flanting opening, entire: Seed 1, within a leaf-like calyx.

H.

lúpulus.

With. Arr. 277.

Flowers greenish yellow. Common Hops. Hedges.

Pen. Di.

XAN THIUM. Male and Fem. flowers on the fame plant.

Male. Cal. common, tiled: Bloff. 1 petal, funnel-shaped, 5-cleft: Recept. chaffy.

Fem. Involucr. 2-leaved, 2-flowered: Bloff. none: Capf. double, prickly, cloven: Nut 2-celled.

Strumárium X. Stem thornless: leaves heartfhaped: 3-fibred.

With. Arr. 283.

Leffer Burdock. Burdock Clotweed. Dunghills. Sept.

Partial Involucr. BUPLEU'RUM. largeft, of 5 leaves: Petals rolled inwards: Fruit egg-shaped, bulging, fmall, not crowned.

rotundifólium.

General involucrum none: B. leaves perforated by the stem.

With. Arr. 285. E. Bot. 99.

Flowers yellowish.

Common Thorough Wax. July. Pen. Di.

> TORDY LIUM. Bloff. radiated, all the florets hermaphrodite: Invo.

lucr. long, undivided: Seeds round, compressed, flat, with a thick border.

T. Umbels crowded, radiated: maximum. leafits spear-shaped, cut, serrated, Bloff. white, red underneath.

With. Arr. 286.

Hartwort. Banks of fields. Pen. Di.

T. Partial involucrums as long as officinále. the flowers: leafits egg-spear-shaped.

With. Arr. 286.

Florets tinged with purple.

Small Hartwort. Corn-fields. Pen. Di.

CAU'CALIS. See Gen. Ch. p. 91.

d g-

\$

9.

y.

all

0.

C. Umbels lateral, fimple, mostly nodos. fitting: feeds egg-shaped, outer ones fet with bristles, inner ones rough.

With. Arr. 289. E. Bot. 199.

Knotted Hen's-foot. Borders of Pen. Di. corn-fields.

Dav'cus. Bloff. fomewhat radiated (generally), all the florets

hermaphrodite: leafits of the involucrum divided: Seeds with membranaceous toothed ridges.

Caróta.

D. Angles of the feeds 4, diftant, hispid: leaf-stalks fibrous underneath: umbel concave when in feed.

With. Arr. 289. Woodv. 161.

Umbels at first flat, or gently convex, but when in seed concave like a tea-cup.

Birds'-neft. Common Wild Carrot. Hedge-fides. Aug.

Pen. Di.

Bu'nium. See Gen. Ch. p. 92.

Bulbocas'-

B. Involucrum of many leaves: frem with leaves at the base rising from the root: styles deciduous.

With, Arr. 291. Curt. 4.

Pen. Di.

Earth Nut.

HERACLE'UM. Involucr. shedding:

Bloff. irregular: Petals bent inwards, notched at the end: Seeds
compressed, leaf-like, smooth,

encompassed by a narrow membranaceous border.

H. Leafits wing-cleft, even: flow- Sphondyers radiated.

With. Arr. 295.

Flowers white.

Cow Parsnep. Madnep. Hedges, July. pastures.

CO'NIUM. Involucellum going half way round, of about 3 leaves: Fruit egg shaped, bulging, ribs compressed, waved before the fruit is ripe.

Seeds without prickles: ftem maculátum. greatly branched, fmooth, fpotted.

With. Arr. 202. Curt. 1. Woodv. 22.

Flowers white.

Common Hemlock. Kex. July. Hedges, orchards, rubbish.

Pen. Di.

SELINUM. Involucr. reflected: Petals heart-shaped, equal: Seeds fmall, with 5 membranaceous ridges, the lateral ones the largest.

S. Stem folitary: ftyles widepalus tre. fpreading after flowering: petals rolled inwards.

With. Arr. 203. E. Bot. 220.

Marsh Milkweed. Wild Parsley. Ditches. July.

Pen. Di.

PEUCED ANUM. Involucrums very fhort: Fruit elliptical, flightly ridged, compreffed and bordered.

P. Leaves 5 times divided into 3: officinále. thread-strap-shaped.

With. Arr. 294.

Common Sulphur-wort. Hogs' Fen-Pen. Di. nel. Salt marshes. July.

minus. P. Leaves winged: leafits wingcleft: fegments strap-shaped, opposite: stem very much branched, fpreading.

Pimpinella. Dioica. Which fee.

With. Arr. 294.

ANGELICA. Bloff. equal: Petals bent inwards: Styles reflected: Fruit roundish.

Pen. Di.

A. Leafits equal, egg-spear-shaped, sylves tris. ferrated.

With. Arr. 298. Woodv. 265.

Bloff. white, tinged with red.

Wild Angelica. Marfhy woods, Pen. Di. hedges. July.

Si'um. Involucrum many - leaved: Petals heart-shaped: Styles bent back: Fruit roundish.

S. Stem creeping: leafits roundish, répens. toothed, and angular.

With. Arr. 300.

Creeping Water Parsnep. July.

Sis on. Involucrum about 4 leaves: Fruit egg-shaped, scored: Styles bent back.

S. Creeping: leaves under water inundatum. hair-like, sthole above winged: umbels of 2 spokes.

With. Arr. 301. E. Bot. 227.

Flowers white.

.

Least Water Parsnep. Water Stonewort. Ditches, pools.

JUNE.

OENAN'THE. Florets of different shapes, those in the centre sitting, barren: Fruit with a cork-like coat, oblong, scored: crowned by the styles and the calyx, which are permanent.

crocáta.

O. All the leaves many-cleft, blunt, nearly equal.

With. Arr. 302. Woodv. 267.

Petals white, acute, bent inwards.

Pen. Di.

Dead Tongue. Hemlock Dropwort. Banks of rivers, ditches. July.

PHELLA'NDRIUM. Central florets fmallest: Fruit egg - shaped, smooth, crowned with the pistil and the calyx.

aquaticum. P. Ramifications of the leaves straddling.

With. Arr. 303. Woodv. 266.

Petals white.

Water Hemlock, or Horsebane.

Pen. Di. Rivers, ditches, pools. July.

Petals bent inwards, notched at the end: Involuctum 1 leaf: Involuctum much production palf-way round: Fruit globular, smooth.

C. Fruit globular. fativ'um, With. Arr. 305. Woodv. 181.

E. Bot. 67.

Flowers whitish.

Coriander. Corn-fields, road- Pen. Di. fides. July.

SCA'NDIX. See Gen. Ch. p. 93.

S. Seeds furrowed, angular. odoráta. With. Arr. 306.

Flowers white.

Sweet Cicely, or Shepherd's Needle.

Orchards, waste places.

Pen. Di.

S. Seeds with a very long beak: Pec'ten. leafits with many fine divisions.

With. Arr. 307. Curt. 5.

Petals white.

Needle. Venus Comb. Corn Pen. Di. fields. July.

IMPERATO'RIA. Petals bent inwards, notched at the end: Seed compressed, with a broad membranaceous border, and 3 ridges on the back.

Ofruthium. I.

With. Arr. 308. Woodv. 35.

Pen. Di. Common Masterwort. Banks of the Clyde.

PASTINA'CA. Petals rolled inwards, entire: Seeds elliptical, compressed, leaf-like, smooth: border thin, narrow.

fativa.

P. Leaves fimply winged.

With. Arr. 309.

Common Wild Parfnep. July.

Pen. Di.

Borders of fields, in limestone.

STAPHYLE'A. Cal. with 5 divisions: Petals 5: Caps. inflated, united: Seeds 2, globular, marked with a fcar, somewhat like a nut.

pinnáta.

S. Leaves fimply winged.

With. Arr. 317.

Bloff. in whirls, white.

Winged Bladder Nut Tree. Hedges near Pontefract.

Pen. Tri.

- CORRIGIO'LA. Cal. 5-leaved, border membranaceous, white: Petals 5: Nut small, roundish, but 3-fided.
- C. Leaves oblong: flowers in a ter- littorális. minating bundle.

With. Arr. 318.

Sand Strapwort. Sands. Sept. Pen. Tri.

LIN'UM. See Gen. Ch. p. 96.

L. Calyx and capfules blunt: leaves peren'ne. fpear-shaped, very entire.

With. Arr. 321. E. Bot. 40.

Bloff. blue.

Perennial Flax.

July.

Pen. Pen.

Pastures in a calcareous soil.

L. Calyx fegments tapering to a tenuifolium. point: leaves fcattered, ftrapawl-shaped, rough when stroked downwards.

With. Arr. 322. E. Bot. 381.

Var. 2. Angustifolium. See E. Bot. p. 381.

Pen. Pen. Narrow-leaved Flax. Dry meadows, pastures. July.

AL'LIUM. See Gen. Ch. p. 98.

arenárium. A. Umbel sheath pointless: stamens 3-pointed: leaf-sheaths cylindrical.

With. Arr. 333.

Bulbs and Bloffoms blue.

Hex. Mo. Sand Garlick. Mountains of Westmoreland.

carinátum. A. Umbel sheath very long: filaments awl-shaped.

With. Arr. 333.

Bloff, dull brown yellow, changing to purple.

Hex. Mo. Mountain Garlick. Rocks, meadows. July.

vinéale. A. Filaments 3-pointed.

With. Arr. 333.

Bloff. small, violet.

Hex. Mo. Crow Garlic. Meadows.

Schænóprafum. A. Leaves cylindrical, awl-threadfhaped, as long as the stalk. With. Arr. 335. Chived Garlick. Meadows, paf-

ORNITHO GALUM. See Gen. Ch. p. 42.

O. Bunch very long: filaments pyrenáicum. fpear-shaped: fruit-stalks when in flower expanding, equal, but afterwards approaching the stalk.

With. Arr. 337.

Petals narrow, expanding, pale yellow.

Spiked Bethlem Star. Meadows, Hex. Mo. paftures.

ANTHE RICUM. Bloff. 6 petals, expanding: Capf. oblong, fuperior, 3-celled.

A. Leaves flattish above, convex scrótinum. underneath: stalk with 1 to 4 flowers: upper floral leaves decurrent, bent inwards.

With. Arr. 339.

Mountain Saffron.

Hex. Mo.

A. Leaves fword-shaped: filaments ofsifragum. woolly.

Narthecium offifragum. Which fee.

Hex. Mo.

With. Arr. 339.

calyculátum. A. Leaves fword-shaped: cup 3lobed: filaments not woolly: flowers with 3 pistils.

Hex. Mo.

Tofielda palustris. Which fee. With. Arr. 339.

Ta'mus. Flowers Male and Fem. on different plants. Cal. with 6 divisions: Bloff. none.

From Style 3-cleft: Berry 3-cel.

Fem. Style 3-cleft: Berry 3-celled, beneath: Seeds 2.

commúnis.

T. Leaves heart-shaped; undivided.

With. Arr. 344. E. Bot. 91.

Bloff. greenish. Berries red.

Bryony Lady-feal. Black Bryony.
Thickets, hedges.

Hex. Mo.

RU'MEX. See Gen. Ch. p. 101.

cris pus.

R. Valves ftrongly veined: leaves fpear-shaped, acute, waved and curled at the edge.

With. Arr. 353. Curt. 2.

Curled Dock. Meadows, road- Hex. Tri. fides. July.

R. Valves veinles: leaves oval- azútus. spear - shaped, uneven at the edges.

With. Arr. 354. Curt. 3.

Woods, hedges. July. Hex. Tri.

R. Leaves oblong, arrow-shaped. Acetosa. With. Arr. 357. E. Bot. 127. Woodv. 60.

Bloff. reddifh.

Common Sorrel. Sorrel Dock.

Meadows, pastures.

Hex. Tri.

ALIS MA. Cal. 3-leaved: Petals 3: Seeds many.

A. Leaves heart-oblong: pistils 6: Damasonium.

With. Arr. 361. Curt. 5.

Bloff. white.

Star-headed Thrunrwort. Aug. Hex. Poly.

A. Leaves strap - spear - shaped, ranuncuon leaf-stalks: fruit globular, loides. scurfy.

With. Arr. 362. E. Bot. 326.

Bloff. bluish white.

Leffer Thrumwort.

Sept.

Hex. Poly.

Wet turfy bogs.

lanceoláta.

A. Leaves fpear-shaped: fruit 3cornered.

With, Arr. 362.

Hex. Poly.

Ditches, shallow waters.

TRIENTA'LIS. Cal. 7 leaves: Bloff. with 7 divisions, equal, flat: Berry dry, superior, globular, 1-valved, 1-celled: Seeds flatted.

europæ'a.

T. Leaves spear-shaped: very entire.

With. Arr. 363. E. Bot. 15.

Bloff. white.

Hept. Mo.

Chickweed Winter Green. July.

EPILO'BIUM. Cal. 4-leaved, deciduous: Petals 4: Capf. beneath, 4-celled, very long: Seeds many, downy at the top.

angustifó-

E. Leaves scattered, strap-spear-shaped: blossoms irregular.

With. Arr. 366. Curt. 2.

Bloff. rose-coloured, or white.

Rosebay Willow Herb. Aug. Oct. Mo.

Watery places.

E. Leaves opposite, egg shaped, montánum. toothed; the upper ones alternate: stem cylindrical.

With. Arr. 367. Curt. 3.

Petals pale purple, fometimes white.

Smooth-leaved Willow Herb.

Woods, hedge-rows. July. Od Mo.

Chlo'ra. Cal. 8-leaved: Bloff. 1 petal, 6 or 8-cleft: Capf. 1-celled, 2 valved, many-feeded.

C. Leaves perforated.

With. Arr. 369. E. Bot. 60.

Bloff. fine full yellow.

Perforated Yellow-wort.
Stiff clay marly grounds. Sept. Oa. Ma.

ERICA. Cal. 4-leaved: Bloff. 4-clefted: Filaments fixed to the receptacle: Anthers cloven and perforated at the top: Capf. 4-celled, 4-valved, many-feeded.

Oft. Mo.

JUNE.

Daboécia. E. Leaves cottony underneath.

With. Arr. 372. E. Bot. 35.

Bloff. purplish red.

Irish Wort, or Moorwort. July. Spongy, wet, uncultivated fand.

vágans. E. Anthers and ftyle projecting: bloffoms bell-shaped: fruit-stalks with 1 flower.

With. Arr. 373. E. Bot. 3.

Oct. Mo. Bloff. dark purple, or rose red.

Aug.

Cornish Heath. Low ground.

Cornish Heath. Low ground.

vulgáris. E. Leaves arrow-shaped.

With. Arr. 374. Curt. 5.

Bloff. pale rose colour, or white.

Oct. Mo. Common Heath. Ling. Heaths, woods. Aug.

fmooth: flowers in umbels, eggfhaped: cups finely ferrated:
branches woolly.

With. Arr. 374. Curt. 2.

Bloff. bluish purple.

Fine-leaved Heath. Dry heaths, Oct. Mo. groves.

Aug.

A'CER. See Gen. Ch. p. 44.

A. Leaves 5-lobed, very entire: campéfire. the lower lobes notched: corymbus with 3 divisions, upright.

With. Arr. 369. E. Bot. 304.

Bloff. terminating, pale green.

Poly'GONUM. See Gen. Ch. p. 45.

P. Style cloven: fpike egg-shaped. amphib'ium With. Arr. 379.

Common Maple. Hedges, thickets. Oa. Mo.

Flowers red. July.

Perennial Willow-leaved Arfmart.

Var. 2. terreftre. Curt. 4.

Leaves darker green.

Amphibious Snakeweed. Sept.

Pools, lakes, marshes, ditches. Off. Tet.

P. Leaves spear shaped: spike ter- vivipárumminating.

With. Arr. 383.

Bloff. whitish.

Oct. Tri.

Small Bistort. Viviparous Snakeweed. Mountains, pastures.

Convol'vu-

P. Leaves heart - shaped: stem twining, angular: flowers blunted.

With. Arr. 385. Curt. 4.

Bloff. greenish white.

Oct, Tri.

Black Bindweed. Climbing Snakeweed. Corn-fields, gardens, hedges. Sept.

RHODIO'LA. Male and Fem. flowers on different plants. Cal. with 4 divisions.

Male. Bloff. 4 petals.

Fem. Bloff. none: Nectaries 4: Capf. 4, many-feeded.

rófea.

R.

With. Arr. 389.

Oct. Tet.

Bloff. terminating, yellow.

Yellow Rosewort. Mountains.

July.

H

Myriophyl'lum. See Gen. Ch. p. 102.

M. Flowers in leafy whirls. verticillá-With. Arr. 390. E. Bot. 218.

Flowers greenish.

Whirled Milfoil. Ditches, Rag- Oct. Tet. nant waters. July.

Bu Tomus. Involucrum fimple, of 3 leaves: Petals 6: Capf. 6, manyfeeded: Seeds fixed to the fides of the capfules.

B. with. Arr. 393. Curt. 1.

Bloff. purple and white.

Flowering Rush. Water Gladiole. Slow streams, muddy ditches. Blo-Norton, Norfolk.

Enne. Hex.

HYDRO'CHARIS. Male and Female flowers on different plants: Cal. 3-cleft: Bloff. 3 petals.

Male. Sheath 2-leaved: Filaments, the 3 inner ones style-bearing.

Fem. Capf. 6-celled, many-feeded, beneath.

H. With. Arr. 394. Curt. 3. Ranz.

Bloff. white.

Common Frogbit. Slow streams, ditches.

Var. 2. Bloffoms doubled, fweetfcented.

Ditch on the fide of Audry Caufey, Enne. Hex. Ifle of Ely.

> MONO TROPA. Cal. none: Petals 10, the five outer ones hollowed at the base to contain honey: Capf. 5-valved. (These are the numbers of the terminating flower.)

Hypop ithys M. Lateral flowers with 8, the terminating flowers with 10 flamens.

> With. Arr. 398. E. Bot. 69. Whole plant fweet, pale yellow. Primrose-scented Birds'-neft.

> > F

Woods, on the roots of other Dec. Mo. July. plants.

> ANDRO MEDA. Cal. with 5 divifions: Bloff. more or less eggfhaped, mouth 5-cleft: Capf. 5celled : Seeds few.

A. Umbel of few flowers, termi- polifólia.

nating: bloffoms egg - fhaped,
nodding: leaves alternate, ftrapfpear-shaped, edges rolled back.

With. Arr. 398.

Bloff. tinged with red.

Marsh Cistus. Wild Rosemary.

Turf bogs.

Dec. Mo.

PY'ROLA. Cal. with 5 divisions:

Petals 5: Caps. 5-celled, opening at the angles: Seeds numerous,
coated.

P. Stamens afcending: piftils lean- rotundifólia. ing downwards.

With. Arr. 400. E. Bot. 213.

Bloff. white.

e

v.

er

y.

i-

g-5Common Winter-green. July.
Woods, groves, in the North. Dec. Mo.

P. Flowers in a bunch, scattered: minor. stamens ascending, pistils straight.

With. Arr. 401. E. Bot. 158.

Bloff. pale pink, or white.

Lesser Winter-green. Woods in Dec. Mothe North. July. fecun da. P. Flowers in a bunch, from one fide.

With. Arr. 401.

Bloff. white.

Dec. Mo. Serrated Winter-green. Woods in the North.

SAXIFRAGA. See Gen. Ch. p. 46.

Relláris.

S. Leaves ferrated, oblong, inversely egg - shaped: stem naked,
branched: petals spear-shaped,
2 dots upon each.

With. Arr. 402. E. Bot. 167.

Petals white.

Hairy Kidney-wort, or Saxifrage.
Moist rocks. July.

and griftly at the edge, rather dented at the end: stem leastless: flowers in a panicle.

With. Arr. 403.

Bloff. whitish, with elegant crimfon dots.

London Pride. None-so-pretty.
Mountains.

Dec. Di.

Dec. Di.

DIAN'THUS. Cal. cylindrical, of 1 leaf, with from 2 to 8 scales at the base: Petals 5, with claws: Caps. cylindrical, 1-celled.

D. Flowers in bundles: scales of barbátus. the calyx egg-awl-shaped, as long as the tube: leaves spear-shaped.

With. Arr. 409.

Sweet Williams. Gardens.

Dec. Di.

D. Scales of the calyx almost rhomboidal, very short: petals irregularly toothed.

Caryophy'llus.

With. Arr. 410. E. Bot. 214. Woody. 80.

Petals pale flesh colour.

Common Pinks. Old walls. July. Dec. Di.

CUCU'BALUS. Calyx inflated: Petals 5, with claws, not crowned at the mouth: Capf. 3-celled.

C. Calyx bell-shaped: petals dif-bacciferus. tant: feed - vessels coloured: branches straddling.

With. Arr. 411.

Flowers white. Berries black.

TUNE.

Dec. Tri. Berry-bearing Chickweed. Hedges in Anglesea. July.

Béhen. C. Calyx nearly globular, smooth, with a net-work of veins: leaves egg - spear - shaped, glaucous, smooth.

With. Arr. 412. E. Bot. 164.

Flowers white.

Spatling Poppy. White Bottle.

Dec. Tri. Corn-fields, dry meadows. Aug.

SILE'NE. See Gen. Ch. p. 48.

an glica. S. Hairy: flowers upright, afternate, on fruit-stalks: lower ones bent back after flowering.

With. Arr. 413. Curt. 4.

Petals white, fometimes with reddish spots.

Dec. Tri. English Catchfly. Corn-fields.
July.

quinquevul nera.

S. Petals very entire, roundish: fruit upright, alternate.

With. Arr. 413. E. Bot. 86.

Dec. Tri. Variegated Catchfly. Sandy cornfields. S. Petals cloven: calyx ribbed: nútans. flowers lateral, pointing one way, bowed downwards: panicle drooping.

With. Arr. 413.

Nottingham Catchfly. Mountain- Dec. Tri. ous meadows. July.

S. Petals cloven: calyx of the cónica. fruit conical, with 30 scores: leaves soft.

With. Arr. 416.

Flowers fine red.

Sandy corn-fields. July. Dec. Tri.

STELLARIA. See Gen. Ch. p. 23.

S. Lower leaves heart-fhaped, on némorum. leaf-stalks: panicle with forked fruit-stalks.

With. Arr. 417. E. Bot. 92.

Flowers white.

Broad-leaved Stichwort. Woods, Dec. Tri.

S. Leaves spear-strap-shaped, very glauca. entire, glaucous green, in cross

pairs: petals half as large again as the calyx.

With. Arr. 420.

Flowers white.

Glaucous -leaved Stichwort.

Dec. Tri. Ditch-banks. July.

uliginófa. S. Leaves spear-shaped, very entire: stem trailing: panicles lateral.

With. Arr. 420. Curt. 6.

Petals white.

Dec. Tri. Pog Stichwort. Sides of springs, boggy meadows.

cerastoídes. S. Leaves oblong: fruit - stalks mostly 2-slowered.

With. Arr. 421.

Petals white.

Dec. Tri. Scotland.

ARENA RIA. See Gen. Ch. p. 104.

peploides. A. Leaves egg - shaped, acute, fleshy.

With. Arr. 421. E. Bot. 189.

Flowers white. July.

Dec. Tri. Sea Sandwort. Sea shore.

A. Leaves thread-shaped, opposite, rúbra. but half the length of the joints of the stem: stem prostrate: calyx as long as the capsule.

With, Arr. 422.

Petals purple.

Purple Spurrey, or Sandwort.
Sandy Meadows.

A

Aug. Dec. Tri.

A. Leaves strap-awl-shaped, 6 in média. a whirl: stem upright: capsules twice the length of the calyx.

With, Arr. 422.

Flowers white.

Pastures on the sea coast. Sept. Dec. Tri.

A. Leaves awl-fhaped: flem pa- tenuifólia. nicled: capfules upright: petals fhorter than the calyx, fpear-fhaped.

With. Arr. 423. E. Bot. 219.

Flowers white.

Fine-leaved Sandwort. Sandy meadows. July. Dec. Tri.

A. Leaves awl-shaped, stiff and juniperina.

fharp: ftems upright: cups fcored: capfules oblong.

With. Arr. 424.

Dec. Tri. Petals white.

COTYLE DON. Cal. 4 or 5-cleft:

Bloff. 1 petal: Nectariferous Scales
5, at the base of the germens:

Caps. 5.

Umbilicus. C. Leaves target or kidney-shaped, fcolloped: stem nearly simple: flowers pendent: floral leaves entire.

With. Arr. 425. E Bot. 325.

Bloff. greenish yellow.

Dec. Pen. Common Navelwort. Kidneywort. Old walls, moift rocks.

Oct.

lútea. C. Leaves target-shaped, scolloped: stem nearly simple: slowers upright: floral leaves toothed.

With. Arr. 426.

Dec. Pen. Yellow Navelwort. Walls, moift rocks.

Se'dum. See Gen. Ch. p. 106.

S. Hairy and viscid: leaves flat-villosum. ted, both them and the leaf-stalks hairy: stem upright, somewhat branched at the base.

With. Arr. 426.

Flowers flesh-coloured.

Marsh Stonecrop.

July. Dec. Pen.

S. Leaves oblong, blunt, nearly al'bum. cylindrical, fitting, expanding: tuft branching.

With. Arr. 427. Curt. 1.

White-flowered Stonecrop. Walls, roofs.

S. Leaves nearly egg-shaped, grow- acre. ing to and fitting, bulging, nearly upright, alternate: tuft with 3 divisions.

With. Arr. 427. Curt. 1, Woodv. 231.

Flowers yellow.

i.

ft

Pepper Stonecrop. Wall Pepper.

Walls, roofs, rocks. July. Dec. Pen.

S. Leaves fomewhat egg-shaped, sexangularegrowing to and sitting, bulglum.

JUNE.

ing, nearly upright, tiled in 6 rows.

With. Arr. 428. Curt. 4.

Flowers yellow.

Dec. Fen. Insipid Stonecrop. Walls, roofs, dry pastures.

S. dafyphyl'-Leaves opposite, egg-shaped, blunt, fleshy: item weak: flowers scattered.

With. Arr. 428. Curt. 3.

Flowers whitish.

Thick-leaved Stonecrop. Dec. Pen. July. Walls, roofs.

> AGROSTEM'MA. Cal. 1 leaf, like leather: Petals 5 with claws: border blunt, undivided: Caps. 1-celled.

Plant hairy: calyx as long as (or longer) than the bloffom: petals entire, not crowned.

With. Arr. 431. Curt. 3.

Flowers purplish red, or white. Corn Cockle. Corn-fields. July.

LYCH'NIS. See Gen. Ch. p. 106.

Githágo.

Dec. Pen.

L. Stamens and pistils in distinct vespertina.
plants: blossoms white: capsules
conical.

With. Arr. 431.

Bloff. white: Capfules conical.

Wild White Campion. Pastures, Dec. Pen. hedges. Aug.

L. Petals 3 or 4 cleft: fruit round- Flos-cúculi. ifh, of 1 cell.

With. Arr. 432. Curt. 1.

Bloff. pink, red, fometimes white.

Meadow Pinks. Ragged Robin.

Var. 2. double flowered.

Pleasant in Sight. Moist mea- Dec. Pen. dows, pastures.

CERA'STIUM. See Gen. Ch. p. 12.

C. Leaves oblong egg - shaped, latifólium. blunt, hairy: branches annual, flaccid, 1 or 2 flowered: capsule oval below, cylindrical, and slightly bent upwards; 10-clest.

With. Arr. 434. Dec. Pen.

C. Leaves oblong, cottony: fruit- tomentofum

JUNE.

flalks branched: capfules globular.

With. Arr. 435.

Petals white.

Dec. Pen. Woolly Mouse-ear. Ripton Wood.

Sper'gula. Cal. 5 leaves: Petals 5, entire: Capf. egg-shaped, of 1 cell and 5 valves.

fubuláta. S. Leaves in whirls, hairy at the edge, rather pointing one way: ftems thread-shaped, flowers solitary.

With. Arr. 436. Curt. 4.

Dec. Pen. Spurry. Aug.

Dry pastures, in gravelly soil.

AGRIMO'NIA. Cal. 5-toothed, furrounded by another: Petals 5: Seeds 2, in a capfule at the bottom of the calyx, which becomes indurated.

Eupatória. A. Stem-leaves winged: the odd leafit on a leaf-stalk: fruit hispid.

With. Arr. 443. Curt. 5.

Woodv. 258.

Flowers fine yellow.

Common Agrimony. Shady places, Dode. Di. hedges. July.

RESE DA. Cal. 1 leaf, divided: Petals jagged: Capf. 1-celled, manyfeeded, opening at the top.

R. Leaves spear - shaped, entire, Lutéola. with a tooth on each side the base: calyx 4-clest.

With. Arr. 445. E. Bot. 320.

Flowers yellow.

Wild Woad. Dyer's Weed. Mea- Dode. Tri. dows, pastures, walls. July.

R. All the leaves 3-cleft: the lútea. lower winged.

With. Arr. 446. E. Bot. 321.

Flowers pale yellow.

Rocket Yellow-weed. Aug.

Meadows, chiefly in a calcare- Dode. Tri. ous foil.

EUPHOR BIA. See Gen. Ch. p. 109.

E. Umbel with 5 spokes; spokes platyphyl'with 3 divisions, and these forkDode. Tri.

ed: involucellums hairy along the keel: leaves ferrated, spearshaped: capsules warty.

With. Arr. 449.

Whole plant yellowish green.

Long-leaved Corn Spurge. Cornfields. July.

Charácias. E. Umbel with many spokes; spokes forked: involucellums perforated, notched at the end: leaves very entire: stem shrublike.

Petals dark purple.

With. Arr. 451.

Dode. Tri. Red Spurge. Woods, hedges.

Spiræ A. Cal. 5-cleft: Petals 5: Capf. 4 or more, 2-celled, 2-valved, many-feeded.

Filipen dula. S. Leaves interruptedly winged:
leafits strap-spear-shaped, irregularly ferrated, very smooth: flowers in tusts.

With. Arr. 463. E. Bot. 284.

Dropwort. Meadowsweet.

Mountainous meadows in a calcareous soil.

July.

Ico. Pen.

S. Leaves interruptedly winged: leafits egg-shaped, doubly serrated, hoary underneath: flowers in tufts.

Ulmária.

With, Arr. 464. Curt. 5.

Flowers yellowish white.

Common Meadowfweet, July.

Moift meadows, river-banks.

Ico. Pen.

Ro'sa. Petals 5: Cal. urn-shaped, 5-cleft, fleshy, contracted at the neck, fo as to form at length a coloured berry of 1 cell, opening at the top: Seeds many, hispid, difperfed in the pulp.

R. Germens and fruit-stalks smooth: spinosis sima ftem and leaf-stalks fully fet with firaight prickles: leafits circular, fmooth.

With. Arr. 465. E. Bot. 187.

Bloff white, cream-colour, or red.

Burnet Rose. Pimpernel Rose. Heaths, fandy places. July. Ico. Poly.

R. Germens and fruit-flalks hispid: villosa. ftem with scattered prickles:

JUNE.

leaf-ftalks prickly: leaves cot-

With. Arr. 466.

Apple Rose. Mountainous hedges.

Var. 2. Wild Briar, or Dog Rose, with large prickly heps.

Ico. Poly. Hedges near London.

rubiginósa. R. Germens and fruit-stalks prickly at the base: prickles on the stem bent back: leaves rusted underneath.

With. Arr. 466.

Sweet Briar Rose. Eglanteria. Hedges, heaths. July.

ftem and leaf-stalks prickly.

With. Arr. 467. Curt. 5.

Woodv. 139.

Petals red, fometimes nearly white.

Dog's Rose. Hep Tree. Wild Briar. Hedges, woods.

Ru'Bus. See Gen. Ch. p. 112.

R. Leaves 3 together, almost bare: lateral leaves 2 -lobed: stem

Ico. Poly.

Ico. Poly.

cæ fius.

prickly, cylindrical: panicle fewflowered.

With. Arr. 468.

Bloff. white. Fruit bluish black. Dewberry Bush, or Bramble.

Woods, hedges. July. Ico. Poly.

R. Leaves winged, with 3 or 5 fruticófus. leafits: stem and leaf-stalks prickly: panicle oblong.

With. Arr. 469.

Framble. Blackberry Bufh.

Hedges. Sept. Ico. Poly.

R. Leaves 3 together, naked: faxatilist fhoots creeping, herbaceous: lower leaves fometimes 5 together.

With. Arr. 470.

Bloff. purple, or white.

Stone Bramble. On the fides of Ico. Poly-

POTENTIL'LA. See Gen. Ch. p. 51.

P. Leaves winged: stem shrub fruiticosa-

With. Arr. 472. E. Bot. 88.

JUNE.

Bloff. yellow.

Shrubby Cinquefoil. South banks of the Tees.

Ico. Poly.

of the rees.

Anserina. P. Leaves winged, serrated: stem creeping: fruit - stalks with 1 flower.

With. Arr. 473. Curt. 3.

Leaves filvery white underneath: Bloff. yellow.

Silver Weed. Wild Tanfey.

Ico. Poly.

Road-fides. July.

fhaped, fnipt, cottony underneath: ftem upright.

With. Arr. 474. E. Bot. 89.

Petals small, yellow.

Silvery, or Hoary Cinquefoil.

Ico. Poly. Meadows in a gravelly foil.

Sept.

rep tans.

P. Leafits 5 together: ftem creeping: fruit-stalks 1-flowered.

With. Arr. 474. Curt. 1: Woodv. 59.

Ico. Poly.

Common Cinquefoil. Meadows, road-fides. Sept.

P. Root-leaves 5-together, serrated, aurea.

pointed: stem-leaves 3 together:

stem declining.

With. Arr. 475. July. Ico. Poly.

Pastures near Pontefract.

TORMENTIL'LA. See Gen. Ch. p. 113.

T. Stem fomewhat ascending: officinális. leaves sitting.

With. Arr. 476. Curt. 5. Woodv. 9.

Septfoil. Tormentil. Moors, paf- Ico. Poly. tures. Sept.

GE'UM. See Gen. Ch. p. 114.

G. Flowers nodding: fruit oblong: rivále. awns feathered, twisted: petals blunt, roundish, wedge-shaped: leaves winged.

With. Arr. 478. E. Bot. 106.

Bloff. ftreaked deadish red.

Water Avens, or Bennet. Sides of rivers, mountains. July. Ico. Poly.

CO'MARUM. Cal. 10-cleft; permanent: fegments alternately fmal-

Poly. Mo.

JUNE

ler: Petals 5: Seeds naked, smooth: Recept. globular, fleshy, woolly, permanent.

palus'tre. C. Leaves winged: petals fmaller than the calyx.

With. Arr. 479. E. Bot. 172.

Ico. Poly. Marsh Cinquefoil. Purple Marsh. locks. Muddy putrid marshes.

July.

PAPA'VER. Bloff. 4 petals: Cal. 2 leaves: Summit target-shaped, radiated, scolloped: Caps. (often) many-celled, opening with holes underneath the permanent summit.

by bridly; frem leafy, many-flowered.

With. Arr. 485. E. Bot. 43.

Petals small, dirty scarlet.

Round Rough-headed Poppy. Cornfields. Norfolk. July.

Argemone. P. Capfule club-shaped, briftly:

With Arr. 486. Curt. 5.

8

Rough - podded Poppy. Corn- Poly. Mo. fields. July.

P. Capfules nearly cylindrical, maritimum. briftly: frem supporting a single flower.

With. Arr. 486. July. Poly. Mo.

P. Capfules smooth, urn-shaped: Rhæ'as. stem hairy, many-flowered: leaves wing-cleft, jagged.

With, Arr. 486. Curt. 3. Woodv. 186.

Bloff. bright scarlet.

Corn, or Red Poppy. Among corn. Aug. Poly. Mo.

P. Capfules oblong, fmooth: ftem dubium. many-flowered: fruit-ftalks with briftles laid to: leaves wing-cleft, fnipt.

With. Arr. 487. Curt. 5.

Smooth-podded Poppy. July. Poly. Mo. Corn-fields in fandy land.

P. Calyx and capfules smooth: somniserum. leaves embracing the stem, jagged.

With. Arr. 487. Woodv. 185.

JUNE.

Poly. Mo. Wild, or White Poppy. Uncultivated gardens. July.

cam bricum. P. Capfules fmooth, oblong: ftem many-flowered, fmooth: leaves winged, jagged.

With. Arr. 488. E. Bot. 66.

Poly. Mo. Yellow Poppy. Mountains in Wales, and about Kendal.

Aug.

Cis'Tus. See Gen. Ch. p. 116.

guttátus. C. Leaves opposite, spear-shaped, 3-sibred: bunches without floral leaves.

With. Arr. 491.

Flowers pale yellow.

Poly. Mo. Annual Ciffus. Sandy pastures, near Holyhead.

bunches, upright: pedicles horizontal.

With. Arr. 491.

Petals yellow.

Poly. Mo. Willow - leaved Ciftus. Sandy meadows. July.

DEEPHIN IUM. Cal. none: Petals
5 or 6: Nectary cloven, hornshaped behind: Capf.leguminous,
many-seeded.

D. Capfule fingle: nectary of 1 Confolida. leaf: ftem sub-divided.

With. Arr. 494.

Flowers blue.

Wild Lark-spier. Corn-fields in Norfolk. Sept. Poly. Tri.

STRATIO TES. Sheath 2 leaves: Cup 3-cleft, or 3 leaves: Petals 3: Berry 6-celled; 6-cornered, beneath.

S. Leaves triangular-Iword-shaped: Aloides. edge fringed with prickles.

With. Arr. 496. E. Bot. 379.

Petals egg-shaped, white.

Water Aloes. Fresh-water Soldier.

Slow streams, fen ditches. July. Poly. Hex.

ZOSTERA. Spike-flalk strap-shaped, concealed within a grass-like leaf, bearing the fructifications on one side: Cal. none: Bloff. none:

Stam. alternate: Capf. alternate: Seeds folitary.

marina. Z. Seed-vessels fitting.

With. Arr. 496.

Poly. Poly. Sea Grafs-wrack. Sea shore.

Aug.

oceánica. Z. Seed-vessels on pedicles, olive-shaped.

With. Arr. 497.

Poly. Poly. Shores of the Isle of Mersey. Aug.

THALIC'TRUM. Cal. none: Petals
4 or 5: Capf. many, rather
beaked.

alpinum. T. Stem unbranched, almost naked: bunch simple, terminating.

With. Arr. 500. E. Bot. 62.

Bloff. brown.

Poly. Poly. Mountain Rue - weed. Moist rocks.

flávum. T. Stem furrowed, leafy: leafits acute, 3-cleft: panicle much branched, upright, compact: flowers upright.

With. Arr. 501. E. Bot. 367.

Flowers yellowish white.

Meadow Rue-weed. Moist mea- Po'y. Poly. dows, river banks.

ADO'NIS. See Gen. Ch. p. 118.

A. Petals about 8: fruit egg-cy- autumnális. lindrical.

With. Arr. 503. Curt. 2. E. Bot. 308.

Petals dark scarlet.

Adonis Flower. Red Morocco.

Corn-fields.

July. Poly. Poly.

RANUN CULUS. See Gen. Ch. p. 7.

R. Leaves egg-spear-shaped, on Flam mula, leaf-stalks: stem declining.

With. Arr. 504. Curt. 6. E. Bot. 387.

Flowers yellow.

Leffer Spearwort. Bogs. Sept. Poly. Poly.

R. Leaves long, spear-shaped: stem Lin'gua. upright.

With. Arr. 504. E. Bot. 100.

Bloff. large, deep yellow.

Great Spearwort, or Crowfoot.

Wet pastures, sides of lakes. Poly. Poly.

Q2 July.

JUNE.

acris.

R. Calyx expanding: fruit-stalks cylindrical: leaves with 3 divifions, and many clefts, the uppermost strap-shaped.

With. Arr. 506. Curt. 1. Woodv. 246. Fl. Ruft. 30.

Bloff yellow.

Butter Flower. Butter Cups.
Meadows, pastures. July.

Poly. Poly.

hederáceus.

R. Leaves roundish, 3-lobed, very entire: stem creeping.

With. Arr. 507. Curt. 4.

Petals spear-shaped, white.

On the mud of flow rivulets.

Poly. Poly.

hirfútus.

R. Root fibrous: stem hairy: calyx pimpled, hairy, taper-pointed, at length reflected.

With. Arr. 508. Curt. 2.

Flowers pale yellow.

Poly. Poly.

Moift clayey places.

Sept.

arven fis.

R. Seeds prickly: upper leaves doubly compound, strap-shaped.

With Arr. 509. Curt. 6.

E. Bot. 135, Fl Rust. 56.

Flowers small, pale yellow. Corn Crowfoot. Corn-fields.

Poly. Poly.

A'juga. See Gen. Ch. p. 54.

A. Leaves downy, fireaked, toothed: floral leaves mostly 3-lobed:
calyxes rough with hair: middle
frem - leaves the largest: rootleaves smaller, tapering into leaf
stalks.

With. Arr. 516.

Mountains in Wales. July. Didy. Gym.

A. Plant woolly, forming a 4- pyramidális. fided pyramid: leaves oblong, toothed: root-leaves largeft.

With, Arr. 516.

Scotland.

Didy. Gym.

VERBE'NA. Bloff. funnel-shaped, fegments nearly equal: Cal. one of its teeth lopped: feeds 2 or 4; naked.

V. Spikes thread-shaped, panicled: officinálisleaves with many jagged clests: stem solitary. Didy. Gym.

Vervain. Simpler's Joy. Waste places, stone walls. Sept. With. Arr. 520. Curt. 1. Woodv. 218.

GALEO'PSIS. Bloff. upper lip vaulted, fomewhat scolloped: lower lip 3-cleft: mouth with a concave taper-pointed tooth on each fide.

Ládanum.

G. All the whirls remote: calyxes bell-shaped, pubescent: teeth longer, strap-bristle shaped, diverging.

With. Arr. 528.

Didy. Gym.

Flowers red. Allheal. Sept. Corn-fields in calcareous foil.

TRU'CRIUM. Upper lip upright, deeply divided, even below the base: stamens in the division.

Chamæ'drys. T. Leaves wedge-egg-shaped, cut, scolloped, on leaf-stalks: flowers together: stems somewhat hairy.

With. Arr. 518. Woodv. 243.

Bloff. reddish purple.

Common Germander. Borders of Didy. Gym. corn-fields. July.

STA'CHYS. Bloff. upper lip vaulted: lower lip bent back at the fides, the larger middle fegment notched: stamens after shedding the pollen bent to the fides.

S. Six flowers in a whirl: leaves fylvatica. heart-shaped, on leaf-stalks.

With. Arr. 531. Curt. 3.

Bloff. deep purple with white fpots.

Var. 2. Leaves angular.

Hedge Nettle Woundwort. Aug. Hedges, woods.

Didy. Gym.

S. Six flowers in a whirl: leaves arven fis. blunt, almost naked: blossoms as long as the calyx: stem feeble.

With. Arr. 532. Curt. 4.

Bloff. whitish, or flesh-coloured.

Aug.

Corn Woundwort. Corn-fields. Didy.Gym.

germánica.

S. Many flowers in a whirl: ferratures of the leaves lapping over each other: flem cottony.

Didy. Gym.

With. Arr. 532. July.

Bloff. purplish red.

Base Horehound. Hedges.

LEONU RUS. Anthers sprinkled with shining particles.

Cardíaca.

L. Stem-leaves spear-shaped, 3-lobed.

With. Arr. 534. E. Bot. 286.

Didy. Gym.

Common Motherwort. Hedges, rubbish, dunghills. Aug.

THY MUS. Cal. 2-lipped: Mouth closed with foft hairs.

ferpyl'lum.

T. Flowers in heads: stems creeping: leaves flat, blunt, fringed at the base.

With. Arr. 536. Curt. 2. Woodw. 110.

Didy. Gym. Mother of Thyme. Wild Thyme. Heaths.

T. Flowers in whirls, 1 upon each A circo.

fruit-flalk: flem upright, fomewhat branched: leaves acute,
ferrated.

With. Arr. 537. Curt. 1.

Bloff. purple.

Wild Bafil. Bafil Thyme. Dry Didy Gym. hills, in chalk and gravel.

Aug.

Melissa. Cal. dry, angular, the upper lip expanding and reflected: Bloff. upper lip fomewhat vaulted, cloven; lower lip, the middle fegment heart shaped.

M. Fruit-stalks axillary, forked, Calamin that as long as the leaves.

With. Arr. 538.

Calamint. Baum. Road fides, corn-fields. Aug. Didy. Gym-

MELITIS. Cal. wider than the tube of the bloffom: Bloff. upper lip upright, entire: lower lip 3-cleft: Anthers, each pair forming a cross.

Melifiophyl'lum. M.

With. Arr. 539. Curt. 6.

Bastard Baum, or Baum-leaf.

Didy. Gym.

Hedges, woods.

PRUNEL'LA. Filaments forked, one of the divisions bearing the anthers: Summit cloven.

vulgáris.

P. All the leaves egg-oblong, ferrated, on leaf-stalks: upper lip of the calyx lopped, 3-toothed.

With. Arr. 541. Curt. 4. Fl. Ruft. 137.

Bloff. blue, purplish, or white.

Didy. Gym.

Common Self-heal. Meadows, pastures. Aug.

RHINAN'THUS. Calyx 4-cleft, inflated: Capfule 2-celled, blunt, compressed: Seeds few.

Crista-gal'li. R. Upper lip of the blossom compressed, and shorter: calyx smooth.

With. Arr. 542. Curt. 5. Fl. Ruft. 138.

Bloff. yellow.

Yellow Rattle. Penny Grafs. Coxcomb. Meadows, woods. July.

Didy. Ang.

MELAMPY RUM. Cal. 4-cleft: Bloff.
upper lip compressed, edges bent
back: Capf. 2-celled, compressed,
slanting, opening on one side:
Seeds solitary, bulging.

M. Spikes quadrangular: floral-criftátum. leaves heart-shaped, compact, finely toothed, tiled.

With. Arr. 544. E. Bot. 41.

Bloff. reddish.

Crefted Cow-wheat. Woods. Didy. Ang. July.

M. Flowers lateral, pointing one fyivat cumway: leaves in diftant pairs: bloffoms widely gaping.

With. Arr. 546.

Yellow Cow-wheat. Aug. Didy. Ang. Woods, shady places.

PEDICULA'RIS. See Gen. Ch. p. 55.

P. Stem branched: calyx crefted alus trise with callous dots: lip of the bloffom flanting.

With. Arr. 548.

Didy. Ang. Bloff. purple, or white. July.

Marst Lousewort. Marshes,
ditches.

Antirrhi num. Cal. with 5 divisions: Bloff. either bulging at the base, or ending in a spur: Caps. 2 - celled, many - seeded, opening at the top, and the divisions bent back.

Cymbalária. A. Leaves heart-shaped, 5-lobed, alternate: stems trailing.

With. Arr. 549. Curt. 1.

Bloff. pale purple.

Ivy-leaved Snapstragon. Oct.

Didy. Ang. Old walls, adjoining the Thames.

fhaped, blunt: stem very much branched, spreading.

With, Arr. 551. Curt. 5.

Didy. Ang. Least Snapdragon. Gravelly cornfields. Sept.

Pelória. A. Var. 4. Bloff. regular, with 5 equal stamens, and 5 nectaries.

With, Arr. 552. Curt. 6.

E. Bot. 260.

Didy. Ang. Sandy fields about Clapham. July.

A. Bloffoms without a fpur: ffow- majus. ers in spikes: cups rounded.

With. Art. 552. E. Bot. 129.

Bloff. rose red, or white.

Greater Snapdragon. Old walls, chalk cliffs. July. Didy. Ang.

LINNÆA. Cal. double, that of the fruit 4-leaved, that of the flower with 5 divisions, superior: Bloff. bell-shaped: Berry dry, 3-celled.

L. Flowers in pairs.

borcális.

With. Arr. 556.

In ancient dry rocky woods.

Didy. Ang.

OROBAN'CHE. See Gen. Ch. p. 122.

O. Stem branched: bloffoms 5- ramófa. cleft.

With. Arr. 558. E. Bot. 184.

Branched Broomrape. Corn-fields, Didy. Ang. dry pastures. Sept:

Bu'nias. Pouch, or rather Drupa, 4-fided, defiduous: angles unequal, acute.

JUNE.

Cákile.

B. Pouch egg-shaped, smooth, with 2 edges and 2 joints.

With, Arr. 562. E. Bot. 231.

Flowers pale purple.

Tetd. Silicu.

Sea Rocket. Sea shore. Oct.

I'satis. Pouch, or Seed-veffel, fimple, oval-spear-shaped, compressed, 1-celled, 1-seeded, crowned by the style.

tinctória.

I. Root -leaves fcolloped: ftemleaves arrow-fhaped: pouches oblong.

With. Arr. 564. E. Bot. 97.

Fl. Ruft, 41.

Tetd. Silicu.

Wild Woad. Corn-fields. July.

Vella. Pouch globular; partition twice as large as the valves, and extending beyond them like the bit of a fpoon.

an ouz.

V. Leaves wing - cleft: pouches pendent.

With. Arr. 564.

Bloff. pale yellow.

Cresse Rocket. Annual Rocket.

Tetd. Silicu.

Salisbury plain.

Subula RIA. Pouch entire, eggfhaped: valves egg fhaped, concave, placed across the partition: Style shorter than the pouch.

S. Leaves awl-shaped. aquática. With. Arr. 564.

Bloff. white.

Water Awlwort. Under water. Tetd.Silicu July.

LEPID'IUM. See Gen. Ch. p. 27.

L. Leaves egg-spear-shaped, en-latifolium. tire, serrated.

With. Arr. 567. E. Bot. 182.

Flowers white.

Dittander Pepperwort. Meadows, Tetd. Silicu. pastures. July.

L. Stem trailing, hairy: leaves did'ymum. with winged clefts: fruit double.

With. Arr. 567. E. Bot. 248.

Flowers white.

Procumbent Dittander. On rub- Tetd. Silicu. bish. July.

L. Root-leaves tooth - winged: ruderále. branch-leaves strap-shaped, very

entire: petals fometimes wanting.

With. Arr. 568.

Narrow-leaved Dittander.

Tetd. Silicu. On rubbish. July.

THLAS PI. See Gen. Ch. p. 26.

T. Pouches round and flat: leaves arven fe. oblong, toothed, fmooth.

With. Arr. 568. Curt. 6.

Treacle Muftard. Penny Cress. Bloff. white. Corn-fields. July.

Tetd. Silicu.

montánum. T. Pouches inverfely-heart-shaped: leaves fmooth: root-leaves battledoor shaped, very entire : stemleaves embracing the ftem: petals longer than the calyx.

With. Arr. 570.

Flowers large, white:

Tetd. Silicu. Pastures in Yorkshire.

COCHUBA RIA. See Gen. Ch. p. 57.

C. Leaves wing-cleft: ftem de-Corénopus, preffed:

With. Art. 574. Fl. Ruft. 92.

Bloff. white.

Swine's Crefs. Scurvy-grafs.

Corn-fields, rubbish. Aug. Tetd. Silicu.

I'BERIS. See Gen. Ch. p. 125.

I. Herbaceous: leaves spear-shap-amára. ed, acute, somewhat toothed: flowers in bunches.

With. Arr. 575. E. Bot. 52.

Flowers white.

Bitter Candy-tuft. Corn-fields. Tetd. Silicu. July.

Sisym'brium. See Gen. Ch. p. 127.

S. Leaves winged: leafits egg- Naftúrtium. shaped.

With. Arr. 580. Curt. 6. Woodv. 48.

Bloff. white.

Common Water-creffes. Springs, Ted. Siliq. brooks. July.

S. Pods oblong-egg-shaped, or cylindrical: leasits spear-shaped, serrated.

With. Arr. 581. Curt. 3.

Flowers small, yellow.

Tetd. Siliq. Creeping Water-crefs. Marshy, watery places. July.

wing cleft, ferrated: petals longer than the cup.

With. Arr. 581.

Flower, the whole yellow.

Tetd. Siliq. Water Radish. Watery places, banks. Aug.

terréstre. S. Root annual: leaves wing-cleft: pods full of seed: petals shorter than the cup.

With. Arr. 582. Curt. 5.

Petals yellow.

Tetd. Siliq. Annual Water-cress. Edges of wet ditches. Sept.

CHEIRAN THUS. See Gen. Ch. p. 128.

what indented: those of the branches entire: pods prickly.

With Arr. 586.

Prickly-podded Gilliflower. Sandy Tetd. Siliq. fea shores. July.

TURRITIS. See Gen. Ch. p. 130.

T. All the leaves hispid: stem- hissúta, leaves embracing the stem.

With. Arr. 589.

Bloff. white.

Wall-crefs. Hairy Towerwort. Rocks, flony places.

Tetd. Siliq.

BRAS'SICA. See Gen. Ch. p. 131.

B. Leaves heart-shaped, embrac- orientális. ing the stem, smooth: root-leaves rough, very entire: pods 4-cornered.

With. Arr. 589.

Petals white.

Perfoliate Cabbage. Cliffs on the Tetd. Siliq. fea coaft.

B. Root and stem steme: stem-campéstris. leaves uniform, heart-shaped, sitting.

With. Arr. 590. Tetd. Siliq.

Flowers yellow, or white.

JUNE.

SINAPIS. See Gen. Ch. p. 132.

nigra.

S. Pods fmooth, laid flat to the fpike-ftalks.

With. Arr 595. Woodv. 151. Fl. Ruft. 51.

Bloff. pale yellow.

Tetd. Siliq.

Common Mustard. Corn-fields, ditch-banks.

RAPHA'NUS. Cal. close, upright:
Necl. glands 2 between the shorter
stamens and the pistil, and 2 between the longer stamens and the
calyx: Pod round, but protuberating, with cells, and nearly
jointed.

Raphinis'-

R. Pods round, jointed, smooth, of 1 cell.

With. Arr. 596. Curt. 4. Fl. Ruft. 71.

Bloff. yellow, or white. July. White-flowered Charlock. Wild Radish. Corn.

Tetd. Siliq.

GERANIUM. See Gen. Ch. p. 59.

G. Petals 2-lobed: leaves circular, pyrenáicum. lower ones with 5, upper with 3 lobes: lobes blunt, 3-cleft,

With. Arr. 601. Curt. 3.

Mountain Crane's-bill. July.

Meadows, Thames bank, Batter- Mond. Dec. fea field.

G. Stem upright: leaves 5-lobed, fylváticum. cut-ferrated: petals entire, or only flightly notched. With. Arr. 602. E Bot. 121.

Petals purple.

Wood Crane's-bill. Mountainous Mond. Dec. thickets. July.

G. Fruit-stalks longer than the columbileaves: leaves 5 - lobed, lobes many-cleft: calyx 5-fided: feedcoats fmooth.

With. Arr. 603. E. Bot. 259.

Long-stalked Crane's-bill. Corn- Mond. Dec. fields, pastures, hedges. Aug.

G. Leaves kidney-shaped, 7 or 5 milvæfolobed: lobes wedge-shaped, 3cleft, bluntish: calyx awnless: feed-coat hairy.

With, Arr. 603, Curt. 6.

Hedges, wafte places.

Var. 1. pufillum. Stem spreading: leaves kidney-shaped, singered: lobes strap-shaped, acute.

Mond. Dec. High ditch banks. July

praten se. G. Leaf - stalks nearly central: leaves with many divisions, wrink-led: lobes with winged clefts, acute: petals entire.

With. Arr. 606. Curt. 4.

Mond. Dec. Crowfoot Crane's-bill. Moissish meadows. July.

maritimum. G. Stem trailing: leaves heartegg-shaped, scolloped, cut, rough: fruit-stalks 1 to 3-flowered: stamens 5.

With, Arr. 606.

Mond. Dec. Sea Crane's-bill. Sea shores.

lúcidum. G. Calyx pyramidal, the angles raised and wrinkled: leaves roundish, 5-lobed.

. With. Arr. 607. E. Bot. 75.

Petals rose red.

Mond. Dec. Shining Crane's-bill. Walls, roofs, rocky places. Aug.

MA'LVA. Cal. double, the outer mostly of 3 leaves: Capf. 8, or more, in a whirl; 1 rarely 2-celled; 1 feed in each.

M. Stem proftrate: leaves heart-rotundifóliafhaped, circular, plaited, 5 or 7lobed: fruit-stalks when ripe declining.

With. Arr. 612. Curt. 3.

Bloff. white, with purple veins.

Dwarf Mallow. Road-fides. Mond. Poly.

M. Stem rough: leaves 5 or 7- sylves bis. lobed, toothed: outer calyx leafits partly united at the base.

With. Arr. 613. Curt. 2.

1th. Arr. 013. Curt. 2. Woody, 54.

Common Mallow. Hedges, foot- Mond. Poly. paths. Aug.

M. Stem upright: root-leaves moscháta. kidney-shaped, cut: stem-leaves with 5 divisions: segments between winged and many-cloven: leasits of the outer cup distinct.

With. Arr. 613. Curt. 4.

Flowers flesh-coloured.

Mond.Poly. Musk Mallow. Meadows, ditchbanks. Aug.

FUMA'RIA. See Gen. Ch. p. 60.

claviculáta. F. Legumens strap-shaped: leaves with tendrils.

With. Arr. 621. E. Bot. 103.

Diad. Hex. Climbing Fumitory. Woods, moist hedges. Sept.

GENISTA. See Gen. Ch. p. 136.

tinctória. G. Branches scored, cylindrical, upright: leaves spear - shaped, smooth: legumen cylindrical.

With. Arr. 624. E. Bot. 44.

Bloff. yellow.

Diad. Dec. Dyer's Greenweed. Pastures, cornfields. Aug.

Ono NIS. Cal. with 5 divisions: Segments strap-shaped: Standard scored: Filaments united without an opening: Legumens swollen, sitting, simple, of 1 cell. O. Flowers in bunches, 2 together: arven'ss. leaves 3-together, the upper ones folitary: branches without thorns, fomewhat woolly.

With. Arr. 627.

Hairy Reft Harrow. Mostly in light land.

Var. 2. Stems proftrate, clammy.

Creeping Reft Harrow. Sea shore, Diad. Dec. in fandy soils.

July.

LA'THYRUS. See Gen. Ch. p. 138.

L. Tendrils without leaves: fti- Aphácapulæ arrow-heart-shaped. With. Arr. 632. Curt. 5.

Flowers yellow.

Yellow Vetchling. Sandy corn Diad. Dec. fields. Sept.

Vi'cia. See Gen. Ch. p. 61.

V. Legumens fitting, bent back, lútea, hairy, folitary: with 5 feeds: standard smooth.

With. Arr. 639.

Yellow Vetch. Meadows. July. Diad. Dec.

hy brida. V. Legumens fitting, bent back, hairy, with 5 feeds: flandard woolly.

With. Arr. 639.

Flowers yellow, greyish above, hairy.

Diad. Dec. Baftard Vetch. Meadows. Aug.

ER'VUM. Cal. with 5 divisions, as long as the blotsom.

tetraspermum. E. Fruit-stalks mostly 2-slowered: feeds 4, globular. With. Arr. 639. Curt. 1.

> Flowers fmall, violet, or bloodcoloured.

Diad. Dec. Smooth-podded Tare. Hedges.

hirsútum. E. Fruit-stalks many-flowered: seeds 2, globular. With. Arr. 640. Curt. 1.

Flowers 2 to 5, pale purple.

Diad. Dec. Wild Tare. Tine Tare. Sandy corn-fields.

Astra Galus. Capf. generally 2celled, bulging; the folitary filaments cylindrical. A. Stem proftrate: flowers in hypoglot'tis, roundish heads: legumens egg-shaped, compressed, hairy, grooved, the point reslected.

With. Arr. 643. E. Bot. 274.

Purple Mountain Milkwort. Diad. Dec. Chalk. July.

A. Stems proftrate: legumens glycyphyl'nearly 3 cornered, bent like a
bow: leaves longer than the fruitftalks: leafits oval.

With. Arr. 643. E. Bot. 203

Flowers greenish yellow.

Wild Liquorice. Meadows, ditch Diad Dec. banks. July.

TRIPO'LIUM. See Gen. Ch. p. 140.

T. Capfules 8-feeded, mostly 3 together, twice as long as the calyx: orn di ftems declining.

Melilótus ornithopodioídes-

With. Arr. 645. Curt. 2.

Bhff. pale red.

Birds'-foot Trefoil. Sandy pastures. Diad. Dec.

Melilótus officinális.

T. Capfules in bunches, often 2feeded, wrinkled, acute: ftem upright.

With. Arr. 645. Fl. Ruft. 72.

Diad. Dec.

Bloff. yellow. Melilot Trefoil. Meadows in stiff foil. July.

ffriatum.

T. Heads fitting, mostly lateral, egg - fhaped: calyxes fcored, rounded.

With. Arr. 649.

Bloff. pale red.

Diad. Dec.

Soft Knotted Trefoil. Dry meadows.

ochroleúcum.

T. Spikes woolly: item upright, pubescent: lower leafits inversely heart-shaped: lower tooth of the calyx as long as the tube of the bloffom.

With. Arr. 653. Fl. Ruft. 35.

Diad. Dec.

Brimftone Trefoil. Dry meadows. July.

maritimum. T. Spikes hairy, globular: ftipulæ fpear - shaped, upright: calyxteeth spreading and dilated after flowering: leafits inverfely eggSpear-shaped: upper leaves opposite.

With. Arr. 653. E. Bot. 220.

Bloff pale reddish purple. July. Diad. Dec. Teafel-headed Trefoil. Sea coaft.

Spikes oval, closely tiled, procumbens many - flowered : ftems widefpreading. With, Arr. 654. Curt. 5.

Flowers upright, yellow.

Hop Trefoil. Dry meadows.

Diad. Dec.

T. Without stem or stalk: flowers suffocátum. nearly fitting on the root.

With. Arr. 656.

Bloff. within the tube of the calyx, colourless.

Yarmouth Denes, near the fea. Diad. Dec.

Calyx tubular: Wings Lo'TUS. converging lengthways and upwards: Legumens straight, generally with cells.

L. Heads of the flowers flatted at corniculatus the top: stems herbaceous, trail-

ing: legumens cylindrical, expanding.

With. Arr. 656. Curt. 2. Fl. Ruft. 53.

Bloff. fweet-fcented, yellow.

Birds'-foot Clover. Aug. Meadows, heaths, road-fides.

Diad. Dec.

Medica'Go. See Gen. Ch. p. 142.

fativa. M. Flowers in bunches: legumens narrow, regular, twisted; stem upright, smooth.

With. Arr. 658. Fl. Ruft. 48.

Bloff. purple.

Lucern Medick. July.

Diad. Dec.

Meadows, ditch-banks.

Hype ricum. Cal. with 5 divisions:

Petals 5: Filaments numerous,
united at the base into 3 or 5 sets:

Capsule with 3 or 5 cells, and
many seeds.

birlutum.

H. Calyx ferrated with glands: frem cylindrical, upright: leaves egg-shaped, formewhat downy.

With. Arr. 666. Curt. 3.

Petals yellow.

Hairy St. John's Wort. Thickets, Polyad.Poly. hedges. July.

H. Calyx ferrated with glands: pul'chrum, frem cylindrical: leaves embracing the stem, heart - shaped, fmooth.

With. Arr. 667. Curt. 1.

Upright St. John's Wort. Hedges, Polyad. Poly. heaths. July.

H. Stem 2-edged: leaves blunt: perforátum. with pellucid dots.

With. Arr. 664. Curt. 1. E. Bot. 205.

Common St. John's Wort. July.

Thickets, hedges, dry banks. Polyad.Poly.

TRAGOPO'GON. See Gen. Ch. p.

T. Calyx as long as the rays of the prater's. blossom: leaves entire, quite straight: fruit-stalk cylindrical.

With. Arr. 672.

Bloff. yellow.

Xellow Goat's-beard. Meadows. Syng. Agua.

Son'chus. Recept. naked: Calyx tiled, bellying: Down hair-like.

eleráceus. S. Fruit - stalks cottony: calyxes

With. Arr. 675. Curt. 2.

Flowers yellow.

Common Sow-thiftle.

Var. 1. Levis. Leaves smooth, the lower with winged clefts, the lower segments long.

Syng, Æqua. Cultivated ground.

Aug.

LEONTO DON. See Gen. Ch. p. 9.

Tarax acum L. Outer scales of the calyx upright, close to the inner, entire: leaves spear-shaped, very entire, or only toothed.

With. Arr. 679. Curt. 1.

Syng. Æqua. Moors, moist places.

hirtum. L. All the scales of the calyx upright: leaves toothed, hairy, hairs undivided: outer row of seeds downless.

With. Arr. 682. Curt. 6.

Flowers yellow.

Syng. Æqua. Heaths, commons. Aug.

HIBRA'CIUM. See Gen. Ch. p. 145.

H. Stem upright, panicle termi- prenanthoinating: leaves spear-shaped, embracing the ftem: glaucous underneath.

With. Arr. 685.

Woods in Scotland.

Syng. Æqua.

- Recept. naked : CRE PIS. double, the outer one deciduous: Down hair-like, fometimes flanding on a pedicle.
- C. Leaves between notched, and foe tida. winged, rough with hair: leafstalks toothed.

With. Arr. 689.

Flowers small, yellow.

Stinking Hawk/beard. Dry mea- Syng. Æqua. dows. July.

C. Leaves spear-shaped, notched, tectorum. fitting, fmooth; the lower ones , toothed.

With. Arr. 689. Curt. 5.

Florets yellow on both fides.

Smooth Succory Hawkfweed. Mea- Syng. Equa. dows, walls, roofs. Sept.

HYPOCHÆRIS. See Gen. Ch. p. 146.

glábra. H. Smooth: stem branched, leafless: leaves toothed and indented: calyx oblong, tiled.

With. Arr. 692. Curt. 3.

Bloff. yellow, close.

Syng. Æqua. , Smooth Cat's-ear. Sandy, gravelly foils. Aug.

LAP'SANA. Recept. naked: Calyx double; all the inner scales channelled.

lar: fruit-stalks slender, very much branched.

With. Arr. 693. Curt. 1.

Bloff. yellow.

Syng. Æqua. Common Nipplewort. Hedges, fhady places. July.

CAR'DUUS. See Gen. Ch. p. 146.

acanthoides. C. Leaves indented, thorny at the edge: calyxes on fruit-stalks, solitary, upright, woolly.

With. Arr. 698.

Flowers purplish, or white. Syng. Equa. Welted Thistle, Road-sides. Sept.

C. Leaves toothed, thorny at the palústris. edge: flowers in bunches, upright: fruit - stalks without thorns.

With. Arr. 697. Curt. 6.

Bloff. purple, or white. July. Syng. Æqua.

C. Leaves decurrent half - way nútans. down towards the next below, thorny: fruit - stalks crooked: calyx scales expanding upwards.

With. Arr. 699.

Flowers fweet - scented in the night.

Musk Thistle. Pastures in a cal-syng. Asqua. careous foil. July.

C. Leaves with winged clefts, hifpid: fegments ftraddling: calyxes egg-shaped, thorny, woolly: ftem hairy.

With. Arr. 699. E. Bot. 107. Fl. Ruft. 131.

Bloff. purple. Aug.

Spear Thiftle. Rubbish, road-sides. Syng. Æqua.

S 2

helenioides

Syng. Æqua.

C, Leaves embracing the stem, spear-shaped, toothed, fringed, with small unequal thorns: stem without thorns.

With. Arr. 702.

Flowers large, purple.

Great English Soft or Melancholy Thistle. Mountainous pastures. July.

CARLI'NA. Cal. radiated: the scales next the blossoms long, coloured.

vulgáris.

C. Stem with many flowers in a corymbus: flowers terminating: rays of the calyx yellow white.

With. Arr. 704.

Syng.Æqua.

Wild Carline Thiftle. Dry meadows.

TANACE TUM. Recept. naked: Down none: Cal. hemispherical, tiled: Florets of the circumference 3clest, narrow-strap-shaped: sometimes wanting.

vulgáre.

T. Leaves double - winged, cut, ferrated.

With. Arr. 708. Woodv. 115.

Flowers in a corymbus, yellow.

Common Tanfy. Banks of rivers. Syng. Super.

SENE CIO. See Gen. Ch. p. 9.

S. Leaves lyre - shaped, almost Jacobæ'a. winged: segments finely jagged: stem upright.

With. Arr. 724. Fl. Ruft. 85.

Bloff. deep yellow.

Ragwort. St. James's Wort. Syng. Super. Meadows, road-fides. July.

CINERA'RIA. See Gen. Ch. p. 147.

C. Flowers in a corymbus: leaves palus tris. broad-fpear-shaped, toothed - indented: stem woolly.

With. Arr. 729. E. Bot. 151.

Bloff. pale yellow.

Marsh Fleawort. Marshes. July. Syng. Super.

CHRYSAN THEMUM. Recept. naked:

Down none, but a kind of border: Cal. hemispherical, tiled;

scales membranaceous at the edge.

Leucan'the
C. Leaves embracing the stem, oblong: ferrated upwards, toothed
at the base.

With. Arr. 733. Curt. 5... Fl. Ruft. 109.

Syng, Super. Greater Daify-Ox-eye. Dry meadows, walls. July.

fégetum. C. Leaves embracing the stem, jagged upwards, tooth-serrated towards the base.

With. Arr. 735. Curt. 6. Fl. Ruft. 110.

Syng.Super. Corn Marigold. Corn-fields. Oct.

MATRICA'RIA. See Gen. Ch. 148.

parthénium. M. Leaves compound, flat: leafits egg-shaped, cut: fruit-stalks branched.

With. Arr. 735. Woodv. 249.

Syng. Super. Common Feverfew. Waste places, hedges, walls. July.

AN'THEMIS. See Gen. Ch. p. 148.

briftle-like: feeds crowned with a border.

With. Arr. 737. Fl. Ruft. 73.

Corn Chamomile. July.

Gravel-pits, corn and fallow- Syng. Super. fields.

ACHILLE'A. Recept. chaffy: Down none: Calyx egg-shaped, tiled: frap-shaped florets from 5 to 10.

A. Leaves strap-spear-shaped, em- Ptarmicabracing the stem, finely serrated. With. Arr. 741. Curt. 5.

Sneezewort. Yarrow. Goofetongue. Moist meadows, shady Syng. Super. places. Aug.

A. Leaves doubly winged, feg- Millefölium. ments of the wings strap-shaped, toothed.

With. Arr. 742. Curt. 6. Woodv. 64. Fl. Ruft. 123.

Bloff. white, or reddish purple.

Yarrow. Millfoil Yarrow. Mea- Syng. Super.

dows, road-fides. Aug.

CENTAURE'A. Recept. bristly: Down either feathered or hair - like: Florets of the circumference funnel-shaped, irregular, longer than the other.

Cy anus,

C. Calyx scales serrated: leaves strap-shaped, very entire: the lower ones toothed.

With. Arr. 742. Curt. 6. E. Bot. 277. Fl. Ruft. 111.

Syng. Frus.

Flue Bottle. Eachelor's Buttons.
Corn-fields. Aug.

Breezewet, Triped Com.

Leave to white wheel,

Twitter to Joilly To

Crements a. Marget hillier Thomas Fither trachered on hell-allies.

A COMPANY OF THE PARTY OF THE P